arXiv:0710.1438v2 [math.PR] 26 Oct 2009

The accuracy of merging approximation in generalized St. Petersburg games
Running title: Accuracy of merging approximation

Gyula Pap
Faculty of Informatics, University of Debrecen,
Pf.12, H–4010 Debrecen, Hungary

April 14, 2009

Abstract

Merging asymptotic expansions of arbitrary length are established for the distribution functions and for the probabilities of suitably centered and normalized cumulative winnings in a full sequence of generalized St. Petersburg games, extending the short expansions due to Csörgő, S., Merging asymptotic expansions in generalized St. Petersburg games, *Acta Sci. Math. (Szeged)* **73** 297–331, 2007. These expansions are given in terms of suitably chosen members from the classes of subsequential semistable infinitely divisible asymptotic distribution functions and certain derivatives of these functions. The length of the expansion depends upon the tail parameter. Both uniform and nonuniform bounds are presented.

AMS 1980 subject classification: Primary 60F05, 60E07; secondary 60G50.

Keywords: merging asymptotic expansions, semistable distribution functions, St. Petersburg games.

1 Introduction

For $p \in (0,1)$ and $\alpha > 0$, consider a generalized St. Petersburg game in which the gain X of a gambler is such that $\mathsf{P}\left\{X = r^{k/\alpha}\right\} = q^{k-1}p$ for $k \in \mathbb{N} := \{1, 2, \dots\}$, where $q := 1-p$ and $r := 1/q$. We refer to Csörgő (4) for a brief introduction to both the history and the mathematics of this game. Let $X_1, X_2, \dots$ be the gambler's gains in a sequence of independent repetitions of the game, with $S_n := X_1 + \cdots + X_n$ standing for the total winnings in n games,

$n \in \mathbb{N}$. The main purpose of the present paper is to study the asymptotic behaviour of the distributions of S_n. Note that $\mathsf{E}(X^\alpha) = \infty$ and $\mathsf{E}(X^\beta) < \infty$ for all $\beta \in (0, \alpha)$, namely,

$$\mathsf{E}(X^\beta) = \frac{p}{q^{\beta/\alpha} - q} =: \mu_\beta^{\alpha,p}.$$

Remark that even for $\beta > \alpha$, a virtual moment $\mu_\beta^{\alpha,p} < 0$ may be defined by this formula. In the finite-variance case $\alpha > 2$, the central limit theorem is valid, namely,

$$\lim_{n\to\infty} \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{a_n^{\alpha,p}} \leq x \right\} = \Phi(x) \qquad \text{for all } x \in \mathbb{R}, \tag{1.1}$$

where $\mathbb{R}$ is the real line, $a_n^{\alpha,p} := \sigma_{\alpha,p}\sqrt{n}$ with $\sigma_{\alpha,p} := \sqrt{\mu_2^{\alpha,p} - [\mu_1^{\alpha,p}]^2}$, and Φ is the standard normal distribution function. In the infinite-variance case $\alpha = 2$ the underlying distribution is still in the domain of attraction of the normal law and (1.1) holds with the choice $a_n^{\alpha,p} := \sqrt{pr\, n \log_r n}$, see Csörgő (4). By standard results for normal approximation, one can derive rate of convergence in the central limit theorem (1.1). Namely, there exist constants $C(\alpha, \beta, p) > 0$ and $C(\alpha, p) > 0$ such that for all $n \in \mathbb{N}$ and $x \in \mathbb{R}$,

$$\left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{a_n^{\alpha,p}} \leq x \right\} - \Phi(x) \right| \leq \begin{cases} \frac{C(\alpha,\beta,p)}{1+|x|^\beta} \frac{1}{\sqrt{n}}, & \text{if } 3 \leq \beta < \alpha, \\ \frac{C(3,p)}{1+x^2} \frac{\log_r n}{\sqrt{n}}, & \text{if } \alpha = 3, \\ \frac{C(\alpha,p)}{1+x^2} \frac{1}{n^{(\alpha-2)/2}}, & \text{if } 2 < \alpha < 3, \\ C(2,p) \frac{1}{\log_r n}, & \text{if } \alpha = 2, \end{cases}$$

see Osipov (16) and Petrov (19, Theorem V.13) for $\alpha > 3$, Hall (11, Theorem 4.1) for $2 < \alpha \leq 3$, Hall (12, Corollary 1) for $\alpha = 2$, and Csörgő (4). Moreover, if $2 < \beta < \alpha \leq 3$ then there exists a bounded decreasing function $\varepsilon_\beta^{\alpha,p} : [1, \infty) \to [0, \infty)$ with $\lim\limits_{u\to\infty} \varepsilon_\beta^{\alpha,p}(u) = 0$ such that for all $n \in \mathbb{N}$ and $x \in \mathbb{R}$,

$$\left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{\sigma_{\alpha,p}\sqrt{n}} \leq x \right\} - \Phi(x) \right| \leq \frac{\varepsilon_\beta^{\alpha,p}\big(\sqrt{n}(1+|x|)\big)}{(1+|x|^\beta)\, n^{(\beta-2)/2}},$$

see Osipov and Petrov (18). The standard results for asymptotic expansion in the central limit theorem (1.1) (see Petrov (19, Theorems VI.1–5)) are not applicable, since the characteristic function of the gain X in one game

$$\mathbf{f}_{\alpha,p}(t) := \mathsf{E}\left(\mathrm{e}^{\mathrm{i}tX}\right) = \sum_{k=1}^{\infty} \mathrm{e}^{\mathrm{i}tr^{k/\alpha}} q^{k-1} p, \qquad t \in \mathbb{R}, \tag{1.2}$$

does not satisfy Cramér's continuity condition $\limsup\limits_{t\to\infty} |\mathbf{f}_{\alpha,p}(t)| < 1$. Indeed, $\mathbf{f}_{\alpha,p}$ is an almost periodic function (see, e.g., Katznelson (13, VI.5)), hence for all $\varepsilon > 0$, there exists $\Lambda > 0$ such that for every $T > 0$, there exists $\tau \in (T, T + \Lambda)$ such that $\sup\limits_{t\in\mathbb{R}} |\mathbf{f}_{\alpha,p}(t) - \mathbf{f}_{\alpha,p}(t + \tau)| < \varepsilon$.

Particularly, $|\mathbf{f}_{\alpha,p}(0)-\mathbf{f}_{\alpha,p}(\tau)|<\varepsilon$, consequently, $\limsup\limits_{t\to\infty}|\mathbf{f}_{\alpha,p}(t)|=1$. However, for $\alpha>3$, in the non-lattice case $r^{1/\alpha}\notin\mathbb{N}$, a short expansion is valid, namely,

$$\sup_{x\in\mathbb{R}}(1+|x|^3)\left|\mathsf{P}\left\{\frac{S_n-\mu_1^{\alpha,p}n}{\sigma_{\alpha,p}\sqrt{n}}\leq x\right\}-\Phi(x)-\frac{Q_1^{\alpha,p}(x)}{\sqrt{n}}\right|=o\left(\frac{1}{\sqrt{n}}\right),$$

as $n\to\infty$ with

$$Q_1^{\alpha,p}(x):=\frac{\mu_3^{\alpha,p}}{6\sqrt{2\pi}[\sigma_{\alpha,p}]^3}(1-x^2)\,\mathrm{e}^{-x^2/2},\qquad x\in\mathbb{R},$$

see Bikelis (2). In the lattice case $r^{1/\alpha}\in\mathbb{N}$, a longer expansion with the usual remainder term can be obtained by adding extra terms to the expansion, see Osipov (17) and Petrov (19, Theorem VI.6).

In the finite-variance lattice case $\alpha>2$ and $r^{1/\alpha}\in\mathbb{N}$, the local limit theorem is also valid, namely,

$$\lim_{n\to\infty}\left[\frac{\sqrt{n}}{r^{1/\alpha}}\,\mathsf{P}\{S_n=s\}-\Phi'_{\alpha,p}\left(\frac{s-\mu_1^{\alpha,p}n}{\sqrt{n}}\right)\right]=0 \tag{1.3}$$

for all $s\in r^{1/\alpha}\mathbb{N}$, where $\Phi_{\alpha,p}(x):=\Phi(x/\sigma_{\alpha,p})$, $x\in\mathbb{R}$. For $\alpha>3$, the rate of convergence is

$$\sup_{s\in r^{1/\alpha}\mathbb{N}}\left|\frac{\sqrt{n}}{r^{1/\alpha}}\,\mathsf{P}\{S_n=s\}-\Phi'_{\alpha,p}\left(\frac{s-\mu_1^{\alpha,p}n}{\sqrt{n}}\right)\right|=O\left(\frac{1}{\sqrt{n}}\right),$$

see Petrov (19, Theorem VII.6). For $2<\alpha\leq 3$, a nonuniform bound

$$\sup_{s\in r^{1/\alpha}\mathbb{N}}\left(1+\left|\frac{s-\mu_1^{\alpha,p}n}{\sqrt{n}}\right|^2\right)\left|\frac{\sqrt{n}}{r^{1/\alpha}}\,\mathsf{P}\{S_n=s\}-\Phi'_{\alpha,p}\left(\frac{s-\mu_1^{\alpha,p}n}{\sqrt{n}}\right)\right|$$
$$=\begin{cases}O\left(\frac{\log_r n}{\sqrt{n}}\right) & \text{if } \alpha=3,\\ O\left(\frac{1}{n^{(\alpha-2)/2}}\right) & \text{if } 2<\alpha<3,\end{cases}$$

is available, see Hall (11, Theorems 5.4 and 5.5). There is also an expansion in the local limit theorem (1.3), namely, for $3\leq\beta<\alpha$,

$$\sup_{s\in r^{1/\alpha}\mathbb{N}}\left(1+\left|\frac{s-\mu_1^{\alpha,p}n}{\sqrt{n}}\right|^{\lfloor\beta\rfloor}\right)\left|\frac{\sqrt{n}}{r^{1/\alpha}}\,\mathsf{P}\{S_n=s\}-\left(U_{\beta,n}^{\alpha,p}\right)'\left(\frac{s-\mu_1^{\alpha,p}n}{\sqrt{n}}\right)\right|=o\left(\frac{1}{n^{(\lfloor\beta\rfloor-2)/2}}\right),$$

where $\lfloor y\rfloor:=\max\{k\in\mathbb{Z}:k\leq y\}$ denotes the lower integer part of a number $y\in\mathbb{R}$, where $\mathbb{Z}:=\{0,\pm1,\pm2,\dots\}$, and

$$U_{\beta,n}^{\alpha,p}(x):=\Phi(x)+\sum_{k=1}^{\lfloor\beta\rfloor-2}\frac{Q_k^{\alpha,p}(x)}{n^{k/2}}$$

is the usual approximating function in the asymptotic expansion in the central limit theorem, where the functions $Q_k^{\alpha,p}$, $k\in\mathbb{N}$, are defined in an appropriate way, see Petrov (19, Theorem VII.16).

From now on, we assume $\alpha \in (0,2)$. Then the distribution function of X is not slowly varying at infinity, hence it is not in the domain of attraction of any (stable) distribution, that is, S_n cannot be centered and normalized to have a proper limit distribution, see Csörgő (4). However, the sequence $(S_n - c_n^{\alpha,p})/n^{1/\alpha}$ has proper limit distributions along some subsequences of $\mathbb{N}$ with centering sequence

$$c_n^{\alpha,p} := \begin{cases} \frac{pn}{q^{1/\alpha}-q} = \mu_1^{\alpha,p} n, & \text{if } \alpha \neq 1, \\ pr\, n \log_r n, & \text{if } \alpha = 1. \end{cases} \tag{1.4}$$

The appropriate subsequences are regulated by the position parameter

$$\gamma_n := \frac{n}{r^{\lceil \log_r n \rceil}} \in (q, 1], \tag{1.5}$$

which describes the location of $n = \gamma_n r^{\lceil \log_r n \rceil}$ between two consecutive powers of $r = 1/q$, where $\lceil y \rceil := \min\{k \in \mathbb{Z} : k \geq y\}$ denotes the upper integer part of a number $y \in \mathbb{R}$. The possible proper limit distributions are semistable infinitely divisible distributions $\{G_{\alpha,p,\gamma} : q < \gamma \leq 1\}$ with exponent α, where $G_{\alpha,p,\gamma}$ can be given by its characteristic function

$$\mathbf{g}_{\alpha,p,\gamma}(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx} \, \mathrm{d}G_{\alpha,p,\gamma}(x) = \mathrm{e}^{y_\gamma^{\alpha,p}(t)}, \qquad t \in \mathbb{R}, \tag{1.6}$$

where

$$y_\gamma^{\alpha,p}(t) := \begin{cases} \sum\limits_{k=-\infty}^{\infty} \left(\exp\left\{ \frac{\mathrm{i}tr^{k/\alpha}}{\gamma^{1/\alpha}} \right\} - 1 \right) \frac{p\gamma}{qr^k}, & \text{if } \alpha \in (0,1), \\ \mathrm{i}tpr \log_r \frac{1}{\gamma} + \sum\limits_{k=-\infty}^{0} \left(\exp\left\{ \frac{\mathrm{i}tr^{k/\alpha}}{\gamma^{1/\alpha}} \right\} - 1 - \frac{\mathrm{i}tr^{k/\alpha}}{\gamma^{1/\alpha}} \right) \frac{p\gamma}{qr^k} & \\ \qquad\qquad + \sum\limits_{k=1}^{\infty} \left(\exp\left\{ \frac{\mathrm{i}tr^{k/\alpha}}{\gamma^{1/\alpha}} \right\} - 1 \right) \frac{p\gamma}{qr^k}, & \text{if } \alpha = 1, \\ \sum\limits_{k=-\infty}^{\infty} \left(\exp\left\{ \frac{\mathrm{i}tr^{k/\alpha}}{\gamma^{1/\alpha}} \right\} - 1 - \frac{\mathrm{i}tr^{k/\alpha}}{\gamma^{1/\alpha}} \right) \frac{p\gamma}{qr^k}, & \text{if } \alpha \in (1,2), \end{cases}$$

for all $t \in \mathbb{R}$, see Csörgő (4) and (6). It can be shown that for any subsequence $\{n_k\}_{k=1}^{\infty}$ of $\mathbb{N}$, the sequence $(S_{n_k} - c_{n_k}^{\alpha,p})/n_k^{1/\alpha}$ converges weakly as $k \to \infty$ if and only if there exists $\gamma \in (q,1]$ such that $\lceil \log_r n_k \rceil - \log_r n_k \to \log_r \frac{1}{\gamma} \pmod 1$ as $k \to \infty$, and in this case

$$\lim_{k \to \infty} \mathsf{P}\left\{ \frac{S_{n_k} - c_{n_k}^{\alpha,p}}{n_k^{1/\alpha}} \leq x \right\} = G_{\alpha,p,\gamma}(x) \qquad \text{for all } x \in \mathbb{R},$$

see Csörgő (8) and Csörgő and Dodunekova (9). It should be noted that the special classical case $\alpha = 1$, $p = 1/2$ and $n_k = 2^k$ has been described by Martin-Löf (15). The above result suggests a merging approximation of the distribution of $(S_n - c_n^{\alpha,p})/n^{1/\alpha}$ by G_{α,p,γ_n}. It turns out that the rates of merge is reasonably fast. Csörgő (4) proved that

$$\sup_{x \in \mathbb{R}} \left| \mathsf{P}\left\{ \frac{S_n - c_n^{\alpha,p}}{n^{1/\alpha}} \leq x \right\} - G_{\alpha,p,\gamma_n}(x) \right| = \begin{cases} O\left(\frac{1}{n}\right), & \text{if } \alpha \in (0,1), \\ O\left(\frac{[\log_r n]^2}{n}\right), & \text{if } \alpha = 1, \\ O\left(\frac{1}{n^{(2-\alpha)/\alpha}}\right), & \text{if } \alpha \in (1,2). \end{cases}$$

In fact, the constants in these bounds are given explicitely by Csörgő (4). For $\alpha \in (0,1]$, Csörgő (8) derived short asymptotic expansions in the merging approximations, namely, for $\alpha \in (0,1)$,

$$\sup_{x\in\mathbb{R}} \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \leq x \right\} - G_{\alpha,p,\gamma_n}(x) + \frac{1}{2n} G_{\alpha,p,\gamma_n}^{(0,2)}(x) \right| = \begin{cases} O\left(\frac{1}{n^2}\right), & \text{if } 0 < \alpha \leq \frac{1}{2}, \\ O\left(\frac{1}{n^{1/\alpha}}\right), & \text{if } \frac{1}{2} < \alpha < 1, \end{cases}$$

for $\alpha = 1$,

$$\sup_{x\in\mathbb{R}} \bigg| \mathsf{P}\left\{ \frac{S_n - pr\, n \log_r n}{n} \leq x \right\}$$
$$- G_{1,p,\gamma_n}(x) - \frac{pr \log_r n}{n} G_{1,p,\gamma_n}^{(1,1)}(x) + \frac{p^2 r^2 [\log_r n]^2}{2n} G_{1,p,\gamma_n}^{(2,0)}(x) \bigg| = O\left(\frac{1}{n}\right),$$

and for $\frac{1}{2} < \alpha < 1$, in the non-lattice case $r^{1/\alpha} \notin \mathbb{N}$,

$$\sup_{x\in\mathbb{R}} \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \leq x \right\} - G_{\alpha,p,\gamma_n}(x) + \frac{1}{2n} G_{\alpha,p,\gamma_n}^{(0,2)}(x) - \frac{\mu_1^{\alpha,p}}{n^{1/\alpha}} G_{\alpha,p,\gamma_n}^{(1,1)}(x) \right| = o\left(\frac{1}{n^{1/\alpha}}\right),$$

where, for $\alpha \in (0,2)$, $p \in (0,1)$, $\gamma \in (q,1]$ and $k, j \in \{0,1,2,\dots\}$, the function $G_{\alpha,p,\gamma}^{(k,j)}$ can be given as a function of bounded variation on the whole $\mathbb{R}$ with Fourier–Stieltjes transform

$$\mathbf{g}_{\alpha,p,\gamma}^{(k,j)}(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\, \mathrm{d}G_{\alpha,p,\gamma}^{(k,j)}(x) = (-\mathrm{i}t)^k \, [y_\gamma^{\alpha,p}(t)]^j \, \mathrm{e}^{y_\gamma^{\alpha,p}(t)}, \qquad t \in \mathbb{R}, \tag{1.7}$$

satisfying $\lim_{x\to-\infty} G_{\alpha,p,\gamma}^{(k,j)}(x) = 0$. The function $G_{\alpha,p,\gamma}^{(k,j)}$ can also be given by

$$G_{\alpha,p,\gamma}^{(k,j)}(x) = \frac{\partial^{k+j} G_{\alpha,p,\gamma}(x,u)}{\partial x^k \, \partial u^j} \bigg|_{u=1}, \qquad x \in \mathbb{R},$$

where, for each $u > 0$, the function $x \mapsto G_{\alpha,p,\gamma}(x,u)$ is a semistable infinitely divisible distribution function given by its characteristic function $t \mapsto \mathbf{g}_{\alpha,p,\gamma}(t,u)$ defined by

$$\mathbf{g}_{\alpha,p,\gamma}(t,u) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\, \mathrm{d}G_{\alpha,p,\gamma}(x,u) = \mathrm{e}^{u\, y_\gamma^{\alpha,p}(t)}, \qquad t \in \mathbb{R},$$

see Csörgő (8).

The aim of the present paper is to establish asymptotic expansions of arbitrary length for $\alpha \in (0,1) \cup (1,2)$, improving the approximation. First, in order to establish candidates for the approximating functions, a formal infinite expansion of the characteristic functions of the cumulative winnings is derived in Section 2, which can be turned into a formal infinite expansion of the distribution functions of the cumulative winnings. Next, in Section 3, a uniform bound is given in case $\alpha \in (0,1) \cup (1,2)$. Its proof, carried out in Section 4, is based on Esseen's classical lemma. In Section 5 a nonuniform bound is presented for $\alpha \in (1,2)$. Its proof in Section 6 is based on a lemma of Osipov (17). The convergence rate of the expansion of length ℓ is $O\left(\frac{1}{n^{1/\alpha}}\right)$ for $\alpha \in \left[\frac{1}{\ell}, 1\right) \cup \left(1, 2-\frac{1}{\ell}\right]$. In the non-lattice case the remainder $O\left(\frac{1}{n^{1/\alpha}}\right)$ will be reduced to $o\left(\frac{1}{n^{1/\alpha}}\right)$. These convergence rates seem to be optimal. The optimality might be studied by the "leading term" approach of Hall (11). In case $\alpha = 1$ one can not improve the short expansion of length 2 obtained by Csörgő (8) by taking a longer expansion, since all the extra terms of a formal infinite expansion are of smaller order than the remainder term $O\left(\frac{1}{n}\right)$. We also give asymptotic expansions of length ℓ in local merging theorems with uniform bound for $\alpha \in (0,1)$ and $\alpha = 1$, and with nonuniform bound for $\alpha \in (1,2)$, with convergence rates $O\left(\frac{1}{n^\ell}\right)$, $O\left(\frac{[\log_r n]^{2\ell}}{n^\ell}\right)$ and $O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right)$, respectively.

Recently Kevei (14) derived uniform merging asymptotic expansions for distribution functions from the domain of geometric partial attraction of a semistable law.

2 Construction of a formal infinite expansion

For fixed $\alpha \in (0,2)$ and $p \in (0,1)$, consider the suitably centered and normalized cumulative winnings $(S_n - c_n^{\alpha,p})/n^{1/\alpha}$. We derive a formal infinite expansion (not taking into consideration its convergence) for the characteristic function

$$\mathbf{f}_n^{\alpha,p}(t) := \mathsf{E}\left(\mathrm{e}^{\mathrm{i}t(S_n - c_n^{\alpha,p})/n^{1/\alpha}}\right), \qquad t \in \mathbb{R},$$

in terms of the Fourier–Stieltjes transforms $\mathbf{g}_{\alpha,p,\gamma_n}^{(k,j)}$, $k, j \in \{0,1,2,\dots\}$. By (1.2),

$$\mathbf{f}_n^{\alpha,p}(t) = \left[\mathbf{f}_{\alpha,p}(t/n^{1/\alpha})\right]^n \mathrm{e}^{-\mathrm{i}tc_n^{\alpha,p}/n^{1/\alpha}} = \left[1 + \frac{x_n^{\alpha,p}(t)}{n}\right]^n \mathrm{e}^{-\mathrm{i}tc_n^{\alpha,p}/n^{1/\alpha}} \tag{2.1}$$

for $t \in \mathbb{R}$, where

$$x_n^{\alpha,p}(t) := n\big(\mathbf{f}_{\alpha,p}(t/n^{1/\alpha}) - 1\big) = n\sum_{k=1}^{\infty}\left(\exp\left\{\frac{\mathrm{i}tr^{k/\alpha}}{n^{1/\alpha}}\right\} - 1\right)q^{k-1}p.$$

It is easy to check that

$$x_n^{\alpha,p}(t) = y_{\gamma_n}^{\alpha,p}(t) + \mathrm{i}t\frac{c_n^{\alpha,p}}{n^{1/\alpha}} + R_{n,1,2}^{\alpha,p}(t), \qquad t \in \mathbb{R}, \tag{2.2}$$

where the position parameter γ_n is given in (1.5), the function $y_\gamma^{\alpha,p}$ stands in the exponent in the definition (1.6) of the characteristic function $\mathbf{g}_{\alpha,p,\gamma}$, and for $k \in \{2,3,\dots\}$ and $t \in \mathbb{R}$,

$$R_{n,1,k}^{\alpha,p}(t) := -\sum_{m=\lceil \log_r n\rceil}^{\infty}\left(\exp\left\{\frac{\mathrm{i}t}{r^{m/\alpha}\gamma_n^{1/\alpha}}\right\} - \sum_{j=0}^{k-1}\frac{(\mathrm{i}t)^j}{j!\,r^{jm/\alpha}\,\gamma_n^{j/\alpha}}\right)\frac{p\gamma_n}{q}r^m, \tag{2.3}$$

see Csörgő (4, page 833). This series is absolutely convergent on the whole line $\mathbb{R}$ for each $k \geq 2$ (see Csörgő (8, page 322) for $k = 2$), thus

$$\begin{aligned}
R_{n,1,k}^{\alpha,p}(t) &= -\frac{p\gamma_n}{q} \sum_{m=\lceil \log_r n \rceil}^{\infty} \sum_{j=k}^{\infty} \frac{(\mathrm{i}t)^j \, r^m}{j! \, r^{jm/\alpha} \, \gamma_n^{j/\alpha}} \\
&= -\frac{p}{q} \sum_{j=k}^{\infty} \frac{(\mathrm{i}t)^j}{j! \, \gamma_n^{(j-\alpha)/\alpha}} \sum_{m=\lceil \log_r n \rceil}^{\infty} q^{m(j-\alpha)/\alpha} \\
&= -\frac{p}{q} \sum_{j=k}^{\infty} \frac{(\mathrm{i}t)^j \, q^{\lceil \log_r n \rceil (j-\alpha)/\alpha}}{j! \, \gamma_n^{(j-\alpha)/\alpha} \, (1 - q^{(j-\alpha)/\alpha})} \\
&= -\frac{p}{q} \sum_{j=k}^{\infty} \frac{(\mathrm{i}t)^j}{j! \, (1 - q^{(j-\alpha)/\alpha}) \, n^{(j-\alpha)/\alpha}} = n \sum_{j=k}^{\infty} \frac{\mu_j^{\alpha,p}}{j!} \left(\frac{\mathrm{i}t}{n^{1/\alpha}} \right)^j . \qquad (2.4)
\end{aligned}$$

Consequently, we have the formal expansion

$$\begin{aligned}
\mathbf{f}_n^{\alpha,p}(t) &= \exp\left\{ -\mathrm{i}t \frac{c_n^{\alpha,p}}{n^{1/\alpha}} + n \log\left(1 + \frac{x_n^{\alpha,p}(t)}{n} \right) \right\} \\
&= \exp\left\{ -\mathrm{i}t \frac{c_n^{\alpha,p}}{n^{1/\alpha}} + x_n^{\alpha,p}(t) + n \sum_{k=2}^{\infty} \frac{(-1)^{k+1}}{k} \left(\frac{x_n^{\alpha,p}(t)}{n} \right)^k \right\} \\
&= \exp\left\{ y_{\gamma_n}^{\alpha,p}(t) + n \sum_{j=2}^{\infty} \frac{\mu_j^{\alpha,p}}{j!} \left(\frac{\mathrm{i}t}{n^{1/\alpha}} \right)^j + n \sum_{k=2}^{\infty} \frac{(-1)^{k+1}}{k} \left(\frac{x_n^{\alpha,p}(t)}{n} \right)^k \right\}.
\end{aligned}$$

First consider the case $\alpha \in (0,1) \cup (1,2)$. Then by (2.2) and (1.4), we have

$$\begin{aligned}
\mathbf{f}_n^{\alpha,p}(t) = \exp\Bigg\{ & y_{\gamma_n}^{\alpha,p}(t) + n \sum_{j=2}^{\infty} \frac{\mu_j^{\alpha,p}}{j!} \left(\frac{\mathrm{i}t}{n^{1/\alpha}} \right)^j + \\
& + n \sum_{k=2}^{\infty} \frac{(-1)^{k+1}}{k} \left[\frac{y_{\gamma_n}^{\alpha,p}(t)}{n} + \sum_{\ell=1}^{\infty} \frac{\mu_\ell^{\alpha,p}}{\ell!} \left(\frac{\mathrm{i}t}{n^{1/\alpha}} \right)^\ell \right]^k \Bigg\} \\
= \mathrm{e}^{y_{\gamma_n}^{\alpha,p}(t)} & \exp\left\{ n \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} d_{k,j}^{\alpha,p} \left(-\frac{\mathrm{i}t}{n^{1/\alpha}} \right)^k \left(\frac{y_{\gamma_n}^{\alpha,p}(t)}{n} \right)^j \right\},
\end{aligned}$$

where the coefficients $d_{k,j}^{\alpha,p}$ are polynomials of the virtual moments $\mu_\ell^{\alpha,p}$, $\ell \in \mathbb{N}$, such that $d_{0,0}^{\alpha,p} = d_{0,1}^{\alpha,p} = d_{1,0}^{\alpha,p} = 0$. Remark that

$$d_{0,2}^{\alpha,p} = -\frac{1}{2}, \qquad d_{1,1}^{\alpha,p} = \mu_1^{\alpha,p}, \qquad d_{2,0}^{\alpha,p} = -\frac{1}{2}[\mu_1^{\alpha,p}]^2 + \frac{1}{2}\mu_2^{\alpha,p},$$

$$d_{0,3}^{\alpha,p} = \frac{1}{3}, \qquad d_{1,2}^{\alpha,p} = -\mu_1^{\alpha,p}, \qquad d_{2,1}^{\alpha,p} = [\mu_1^{\alpha,p}]^2 - \frac{1}{2}\mu_2^{\alpha,p}, d_{3,0}^{\alpha,p} = -\frac{1}{3}[\mu_1^{\alpha,p}]^3 + \frac{1}{2}\mu_1^{\alpha,p}\mu_2^{\alpha,p} - \frac{1}{6}\mu_3^{\alpha,p},$$

$$d_{0,4}^{\alpha,p} = -\frac{1}{4}, \qquad d_{1,3}^{\alpha,p} = \mu_1^{\alpha,p}, \qquad d_{2,2}^{\alpha,p} = -\frac{3}{2}[\mu_1^{\alpha,p}]^2 + \frac{1}{2}\mu_2^{\alpha,p}, \qquad d_{3,1}^{\alpha,p} = [\mu_1^{\alpha,p}]^3 - \mu_1^{\alpha,p}\mu_2^{\alpha,p} + \frac{1}{6}\mu_3^{\alpha,p},$$

$$d_{4,0}^{\alpha,p} = -\frac{1}{4}[\mu_1^{\alpha,p}]^4 + \frac{1}{2}[\mu_1^{\alpha,p}]^2\mu_2^{\alpha,p} - \frac{1}{6}\mu_1^{\alpha,p}\mu_3^{\alpha,p} - \frac{1}{8}[\mu_2^{\alpha,p}]^2 + \frac{1}{24}\mu_4^{\alpha,p}.$$

Following Christoph and Wolf (3, Section 4.3), using the Taylor expansion of the exponential function, we obtain a formal power series expansion

$$\mathbf{f}_n^{\alpha,p}(t) = \mathrm{e}^{y_{\gamma_n}^{\alpha,p}(t)}\left\{1 + \sum_{k=0}^{\infty}\sum_{j=-\lfloor k/2\rfloor}^{\infty}\sum_{m=\max\{1,-j\}}^{k+j}\frac{w_{m,k,j}^{\alpha,p}\,(-\mathrm{i}t)^k\,[y_{\gamma_n}^{\alpha,p}(t)]^{j+m}}{m!\,n^{\frac{k}{\alpha}+j}}\right\},$$

where

$$w_{m,k,j}^{\alpha,p} := \begin{cases} 0, & \text{if } -\lfloor k/2\rfloor \le j < L_{m,k}, \\ \displaystyle\sideset{}{'}\sum_{\substack{k_1+\cdots+k_m=k\\ s_1+\cdots+s_m=j+m}} d_{k_1,s_1}^{\alpha,p}\ldots d_{k_m,s_m}^{\alpha,p}, & \text{if } j \ge L_{m,k}, \end{cases}$$

with

$$L_{m,k} := \max\{-\lfloor k/2\rfloor, -m, m-k\}.$$

The summation $\sum'$ in the definition of $w_{m,k,j}^{\alpha,p}$ is carried over all nonnegative integers $k_1, \ldots, k_m, s_1, \ldots, s_m$, such that $k_1 + \cdots + k_m = k$, $s_1 + \cdots + s_m = j + m$, and $(k_j, s_j) \notin \{(0,0),(1,0),(0,1)\}$. Finally, we conclude a formal expansion

$$\mathbf{f}_n^{\alpha,p}(t) = \mathbf{g}_{\alpha,p,\gamma_n}(t) + \sum_{k=0}^{\infty}\sum_{j=-\lfloor k/2\rfloor}^{\infty}\sum_{m=\max\{1,-j\}}^{k+j}\frac{w_{m,k,j}^{\alpha,p}\,\mathbf{g}_{\alpha,p,\gamma_n}^{(k,\,j+m)}(t)}{m!\,n^{\frac{k}{\alpha}+j}}, \tag{2.5}$$

where the position parameter γ_n is introduced in (1.5), the characteristic function $\mathbf{g}_{\alpha,p,\gamma}$ is defined in (1.6), and the function $\mathbf{g}_{\alpha,p,\gamma}^{(k,\,j)}$ is given in (1.7). Replacing each term by its inverse Fourier-Stieltjes transform, this may be turned into a formal expansion

$$\mathsf{P}\left\{\frac{S_n - \mu_1^{\alpha,p}n}{n^{1/\alpha}} \le x\right\} = G_{\alpha,p,\gamma_n}(x) + \sum_{k=0}^{\infty}\sum_{j=-\lfloor k/2\rfloor}^{\infty}\sum_{m=\max\{1,-j\}}^{k+j}\frac{w_{m,k,j}^{\alpha,p}\,G_{\alpha,p,\gamma_n}^{(k,\,j+m)}(x)}{m!\,n^{\frac{k}{\alpha}+j}},$$

for $x \in \mathbb{R}$.

Now consider the case $\alpha = 1$. Then by (2.2) and (1.4), we have

$$\begin{aligned}\mathbf{f}_n^{1,p}(t) &= \exp\left\{ y_{\gamma_n}^{1,p}(t) + n\sum_{j=2}^{\infty} \frac{\mu_j^{1,p}}{j!}\left(\frac{\mathrm{i}t}{n}\right)^j \right.\\ &\qquad \left. + n\sum_{k=2}^{\infty}\frac{(-1)^{k+1}}{k}\left[\frac{y_{\gamma_n}^{1,p}(t) + \mathrm{i}tpr\log_r n}{n} + \sum_{\ell=2}^{\infty}\frac{\mu_\ell^{1,p}}{\ell!}\left(\frac{\mathrm{i}t}{n}\right)^\ell\right]^k \right\}\\ &= \mathrm{e}^{y_{\gamma_n}^{1,p}(t)}\exp\left\{ n\sum_{k=0}^{\infty}\sum_{j=0}^{\infty} d_{k,j}^{1,p}\left(-\frac{\mathrm{i}t}{n}\right)^k\left(\frac{y_{\gamma_n}^{1,p}(t) + \mathrm{i}tpr\log_r n}{n}\right)^j \right\},\end{aligned}$$

where the coefficients $d_{k,j}^{1,p}$ are now polynomials of the virtual moments $\mu_\ell^{1,p}$ with only $\ell \geq 2$, such that $d_{0,0}^{1,p} = d_{0,1}^{1,p} = d_{1,0}^{1,p} = 0$. Remark that a formula for $d_{k,j}^{1,p}$ can be obtained from the formula for $d_{k,j}^{\alpha,p}$, $\alpha \in (0,1)\cup(1,2)$, by replacing α by 1 and $\mu_1^{1,p}$ by 0. Hence following Christoph and Wolf (3, Section 4.3), now we obtain a formal power series expansion

$$\mathrm{e}^{y_{\gamma_n}^{1,p}(t)}\left\{1 + \sum_{k=0}^{\infty}\sum_{j=-\lfloor k/2\rfloor}^{\infty}\sum_{m=\max\{1,-j\}}^{k+j}\frac{w_{m,k,j}^{1,p}\,(-\mathrm{i}t)^k\,[y_{\gamma_n}^{1,p}(t) + \mathrm{i}tpr\log_r n]^{j+m}}{m!\,n^{k+j}}\right\}$$

for $\mathbf{f}_n^{1,p}(t)$, $t \in \mathbb{R}$, which may be turned into a formal expansion

$$\begin{aligned}\mathsf{P}\left\{\frac{S_n - pr\,n\log_r n}{n} \leq x\right\} &= G_{1,p,\gamma_n}(x) + \sum_{k=0}^{\infty}\sum_{j=-\lfloor k/2\rfloor}^{\infty}\sum_{m=\max\{1,-j\}}^{k+j}\frac{w_{m,k,j}^{1,p}}{m!\,n^{k+j}}\\ &\qquad\times\sum_{\ell=0}^{j+m}\binom{j+m}{\ell}[-pr\log_r n]^\ell\,G_{1,p,\gamma_n}^{(k+\ell,\,j+m-\ell)}(x),\end{aligned}$$

for $x \in \mathbb{R}$. On the other hand, the above formula for $\mathbf{f}_n^{1,p}(t)$, and hence the above expansion for $\mathsf{P}\left\{\frac{S_n - pr\,n\log_r n}{n} \leq x\right\}$ in case $\alpha = 1$ can be obtained from the formula for $\mathbf{f}_n^{\alpha,p}(t)$, $\alpha \in (0,1)\cup(1,2)$, and from the expansion for $\mathsf{P}\left\{\frac{S_n - \mu_1^{\alpha,p}n}{n^{1/\alpha}} \leq x\right\}$ by replacing α by 1 and $\mu_1^{1,p}$ by $pr\log_r n$.

3 Uniform bounds in asymptotic expansions

For $\alpha \in (0,1)\cup(1,2)$, $p \in (0,1)$, $n \in \mathbb{N}$ and $\ell \in \{0,1,\dots\}$, introduce

$$G_{n,\ell}^{\alpha,p}(x) := G_{\alpha,p,\gamma_n}(x) + \sum_{k=0}^{2\ell}\sum_{j=-\lfloor k/2\rfloor}^{\ell-k}\sum_{m=\max\{1,-j\}}^{k+j}\frac{w_{m,k,j}^{\alpha,p}\,G_{\alpha,p,\gamma_n}^{(k,j+m)}(x)}{m!\,n^{\frac{k}{\alpha}+j}}$$

for $x \in \mathbb{R}$. The function $G^{\alpha,p}_{n,\ell}$ consists of all terms of order $n^{-\frac{k}{\alpha}-j}$ with $k+j \le \ell$ of the formal expansion. Remark that $G^{\alpha,p}_{n,0} = G_{\alpha,p,\gamma_n}$, and

$$
\begin{aligned}
G^{\alpha,p}_{n,1} &= G_{\alpha,p,\gamma_n} + \frac{d^{\alpha,p}_{0,2}\, G^{(0,2)}_{\alpha,p,\gamma_n}}{n} + \frac{d^{\alpha,p}_{1,1}\, G^{(1,1)}_{\alpha,p,\gamma_n}}{n^{\frac{1}{\alpha}}} + \frac{d^{\alpha,p}_{2,0}\, G^{(2,0)}_{\alpha,p,\gamma_n}}{n^{\frac{2}{\alpha}-1}} \\
&= G_{\alpha,p,\gamma_n} - \frac{G^{(0,2)}_{\alpha,p,\gamma_n}}{2n} + \frac{\mu^{\alpha,p}_1\, G^{(1,1)}_{\alpha,p,\gamma_n}}{n^{\frac{1}{\alpha}}} - \frac{\big([\mu^{\alpha,p}_1]^2 - \mu^{\alpha,p}_2\big)\, G^{(2,0)}_{\alpha,p,\gamma_n}}{2n^{\frac{2}{\alpha}-1}},
\end{aligned}
$$

$$
\begin{aligned}
G^{\alpha,p}_{n,2} = G^{\alpha,p}_{n,1} &+ \frac{d^{\alpha,p}_{0,3}\, G^{(0,3)}_{\alpha,p,\gamma_n} + \frac{1}{2}[d^{\alpha,p}_{0,2}]^2\, G^{(0,4)}_{\alpha,p,\gamma_n}}{n^2} + \frac{d^{\alpha,p}_{1,2}\, G^{(1,2)}_{\alpha,p,\gamma_n} + d^{\alpha,p}_{0,2}\, d^{\alpha,p}_{1,1}\, G^{(1,3)}_{\alpha,p,\gamma_n}}{n^{1+\frac{1}{\alpha}}} \\
&+ \frac{d^{\alpha,p}_{2,1}\, G^{(2,1)}_{\alpha,p,\gamma_n} + \frac{1}{2}\big([d^{\alpha,p}_{1,1}]^2 + 2d^{\alpha,p}_{2,0} d^{\alpha,p}_{0,2}\big)\, G^{(2,2)}_{\alpha,p,\gamma_n}}{n^{\frac{2}{\alpha}}} \\
&+ \frac{d^{\alpha,p}_{3,0}\, G^{(3,0)}_{\alpha,p,\gamma_n} + d^{\alpha,p}_{1,1}\, d^{\alpha,p}_{2,0}\, G^{(3,1)}_{\alpha,p,\gamma_n}}{n^{\frac{3}{\alpha}-1}} + \frac{\frac{1}{2}[d^{\alpha,p}_{2,0}]^2\, G^{(4,0)}_{\alpha,p,\gamma_n}}{n^{\frac{4}{\alpha}-2}} \\
= G^{\alpha,p}_{n,1} &+ \frac{8G^{(0,3)}_{\alpha,p,\gamma_n} + 3G^{(0,4)}_{\alpha,p,\gamma_n}}{24\, n^2} - \frac{\mu^{\alpha,p}_1\big(2G^{(1,2)}_{\alpha,p,\gamma_n} + G^{(1,3)}_{\alpha,p,\gamma_n}\big)}{2n^{1+\frac{1}{\alpha}}} \\
&+ \frac{2\big(2[\mu^{\alpha,p}_1]^2 - \mu^{\alpha,p}_2\big)\, G^{(2,1)}_{\alpha,p,\gamma_n} + \big(3[\mu^{\alpha,p}_1]^2 - \mu^{\alpha,p}_2\big)\, G^{(2,2)}_{\alpha,p,\gamma_n}}{4n^{\frac{2}{\alpha}}} \\
&- \frac{\big(2[\mu^{\alpha,p}_1]^3 - 3\mu^{\alpha,p}_1\, \mu^{\alpha,p}_2 + \mu^{\alpha,p}_3\big)\, G^{(3,0)}_{\alpha,p,\gamma_n} + 3\mu^{\alpha,p}_1\big([\mu^{\alpha,p}_1]^2 - \mu^{\alpha,p}_2\big)\, G^{(3,1)}_{\alpha,p,\gamma_n}}{6n^{\frac{3}{\alpha}-1}} \\
&+ \frac{\big([\mu^{\alpha,p}_1]^2 - \mu^{\alpha,p}_2\big)^2\, G^{(4,0)}_{\alpha,p,\gamma_n}}{8n^{\frac{4}{\alpha}-2}}.
\end{aligned}
$$

For $\alpha = 1$, $p \in (0,1)$, $n \in \mathbb{N}$ and $\ell \in \{0,1,\dots\}$, introduce

$$
\begin{aligned}
G^{1,p}_{n,\ell}(x) := G_{1,p,\gamma_n}(x) + \sum_{k=0}^{2\ell} \sum_{j=-\lfloor k/2 \rfloor}^{\ell-k} \sum_{m=\max\{1,-j\}}^{k+j} \frac{w^{1,p}_{m,k,j}}{m!\, n^{k+j}} \\
\times \sum_{\ell=0}^{j+m} \binom{j+m}{\ell} [-pr \log_r n]^{\ell}\, G^{(k+\ell,\, j+m-\ell)}_{1,p,\gamma_n}(x),
\end{aligned}
$$

for $x \in \mathbb{R}$. The function $G^{1,p}_{n,\ell}$ consists of all terms containing n^{-k-j} with $k+j \le \ell$ of the formal expansion. Remark that $G^{1,p}_{n,0} = G_{1,p,\gamma_n}$, and

$$
G^{1,p}_{n,1} = G_{1,p,\gamma_n} - \frac{G^{(0,2)}_{1,p,\gamma_n}}{2n} + \frac{[pr \log_r n]\, G^{(1,1)}_{1,p,\gamma_n}}{n} - \frac{\big([pr \log_r n]^2 - \mu^{1,p}_2\big)\, G^{(2,0)}_{1,p,\gamma_n}}{2n},
$$

$$\begin{aligned}G^{1,p}_{n,2} = G^{1,p}_{n,1} &+ \frac{8G^{(0,3)}_{1,p,\gamma_n} + 3G^{(0,4)}_{1,p,\gamma_n}}{24\,n^2} - \frac{[pr\log_r n]\bigl(2G^{(1,2)}_{1,p,\gamma_n} + G^{(1,3)}_{1,p,\gamma_n}\bigr)}{2n^2}\\ &+ \frac{2\bigl(2[pr\log_r n]^2 - \mu^{1,p}_2\bigr)\,G^{(2,1)}_{1,p,\gamma_n} + \bigl(3[pr\log_r n]^2 - \mu^{1,p}_2\bigr)\,G^{(2,2)}_{1,p,\gamma_n}}{4n^2}\\ &- \frac{\bigl(2[pr\log_r n]^3 - 3[pr\log_r n]\,\mu^{1,p}_2 + \mu^{1,p}_3\bigr)\,G^{(3,0)}_{1,p,\gamma_n}}{6n^2}\\ &- \frac{3[pr\log_r n]\bigl([pr\log_r n]^2 - \mu^{1,p}_2\bigr)\,G^{(3,1)}_{1,p,\gamma_n}}{6n^2} + \frac{\bigl([pr\log_r n]^2 - \mu^{1,p}_2\bigr)^2\,G^{(4,0)}_{1,p,\gamma_n}}{8n^2}.\end{aligned}$$

The functions $G^{\alpha,p}_{n,1}$, $\alpha \in (0,2)$, are almost the same as the approximating functions $G^{\alpha,p}_{n}$, $\alpha \in (0,2)$, of Csörgő (8). They differ only in the coefficients of their last terms, which does not make difference in the convergence rates if $\alpha \in (0,1]$, but the functions $G^{\alpha,p}_{n,1}$, $\alpha \in (1,2)$, give better approximations than $G^{\alpha,p}_{n}$, $\alpha \in (1,2)$, of Csörgő (8).

The main results of this section are contained in the following

Proposition 3.1. *For* $\ell \in \mathbb{N}$,

$$\sup_{x\in\mathbb{R}}\left|\mathsf{P}\left\{\frac{S_n - \mu^{\alpha,p}_1 n}{n^{1/\alpha}} \le x\right\} - G^{\alpha,p}_{n,\ell-1}(x)\right| = \begin{cases} O\bigl(\frac{1}{n^\ell}\bigr)\,, & \textit{if } \ 0 < \alpha < \frac{1}{\ell},\\ O\bigl(\frac{1}{n^{1/\alpha}}\bigr)\,, & \textit{if } \ \frac{1}{\ell} \le \alpha < 1 \ \ \textit{or} \ \ 1 < \alpha \le 2 - \frac{1}{\ell},\\ O\bigl(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\bigr)\,, & \textit{if } \ 2 - \frac{1}{\ell} < \alpha < 2.\end{cases}$$

For $\ell \in \{2,3,\dots\}$, $\alpha \in \bigl(\frac{1}{\ell},1\bigr) \cup \bigl(1, 2-\frac{1}{\ell}\bigr)$ *and* $r^{1/\alpha} \notin \mathbb{N}$,

$$\sup_{x\in\mathbb{R}}\left|\mathsf{P}\left\{\frac{S_n - \mu^{\alpha,p}_1 n}{n^{1/\alpha}} \le x\right\} - G^{\alpha,p}_{n,\ell-1}(x)\right| = o\left(\frac{1}{n^{1/\alpha}}\right).$$

In case $\ell = 1$, the approximating function is $G^{\alpha,p}_{n,0} = G_{\alpha,p,\gamma_n}$, and Proposition 3.1 gives back the rates of merge for $0 < \alpha < 1$ and $1 < \alpha < 2$, due to Csörgő (4). Certain terms of $G^{\alpha,p}_{n,\ell-1}$ are of the same or of a smaller order than the remainder terms $O\bigl(\frac{1}{n^{1/\alpha}}\bigr)$ or $o\bigl(\frac{1}{n^{1/\alpha}}\bigr)$. Using boundedness of the functions $G^{(k,j)}_{\alpha,p,\gamma_n}$ (see Csörgő (8, Lemma 6)), the expansions may be simplified as follows.

Theorem 3.1. *For* $\ell \in \{2,3,\dots\}$, $\alpha \in \bigl[\frac{1}{\ell},\frac{1}{\ell-1}\bigr) \cup \bigl(2-\frac{1}{\ell-1},\, 2-\frac{1}{\ell}\bigr]$ *and* $r^{1/\alpha} \in \mathbb{N}$,

$$\sup_{x\in\mathbb{R}}\left|\mathsf{P}\left\{\frac{S_n - \mu^{\alpha,p}_1 n}{n^{1/\alpha}} \le x\right\} - \tilde{G}^{\alpha,p}_{n,\ell-1}(x)\right| = O\left(\frac{1}{n^{1/\alpha}}\right);$$

for $\ell = 2$, $\alpha \in \bigl(\frac{1}{2},1\bigr) \cup \bigl(1,\frac{3}{2}\bigr)$, *or for* $\ell \in \{3,4,\dots\}$, $\alpha \in \bigl(\frac{1}{\ell},\frac{1}{\ell-1}\bigr] \cup \bigl[2-\frac{1}{\ell-1},\, 2-\frac{1}{\ell}\bigr)$, *and for* $r^{1/\alpha} \notin \mathbb{N}$,

$$\sup_{x\in\mathbb{R}}\left|\mathsf{P}\left\{\frac{S_n - \mu^{\alpha,p}_1 n}{n^{1/\alpha}} \le x\right\} - \tilde{\tilde{G}}^{\alpha,p}_{n,\ell-1}(x)\right| = o\left(\frac{1}{n^{1/\alpha}}\right),$$

where the approximating functions are given by

$$\tilde{G}^{\alpha,p}_{n,\ell-1} := \begin{cases} G_{\alpha,p,\gamma_n} + \displaystyle\sum_{j=1}^{\ell-1}\sum_{m=1}^{j} \frac{w^{\alpha,p}_{m,0,j}\, G^{(0,j+m)}_{\alpha,p,\gamma_n}}{m!\, n^j}, & \textit{if} \ \ \alpha \in (0,1), \\ G_{\alpha,p,\gamma_n} + \displaystyle\sum_{j=1}^{\ell-1} \frac{(-1)^j \big([\mu^{\alpha,p}_1]^2 - \mu^{\alpha,p}_2\big)^j\, G^{(2j,0)}_{\alpha,p,\gamma_n}}{m!\, 2^j\, n^{j(2-\alpha)/\alpha}}, & \textit{if} \ \ \alpha \in (1,2), \end{cases}$$

and $\tilde{\tilde{G}}^{\alpha,p}_{n,\ell-1} := \tilde{G}^{\alpha,p}_{n,\ell-1} + \dfrac{\mu^{\alpha,p}_1\, G^{(1,1)}_{\alpha,p,\gamma_n}}{n^{1/\alpha}}$.

If $\ell = 2$ then the approximating function is

$$\tilde{G}^{\alpha,p}_{n,1} = \begin{cases} G_{\alpha,p,\gamma_n} - \dfrac{G^{(0,2)}_{\alpha,p,\gamma_n}}{2n}, & \text{if} \ \ \alpha \in (0,1), \\ G_{\alpha,p,\gamma_n} - \dfrac{\big([\mu^{\alpha,p}_1]^2 - \mu^{\alpha,p}_2\big)\, G^{(2,0)}_{\alpha,p,\gamma_n}}{2\, n^{(2-\alpha)/\alpha}}, & \text{if} \ \ \alpha \in (1,2), \end{cases}$$

and for $\alpha \in (0,1)$ the result has been obtained by Csörgő (8). The approximating functions for $\ell = 3$, $\ell = 4$ and $\alpha \in (0,1)$ are

$$\tilde{G}^{\alpha,p}_{n,2} = \tilde{G}^{\alpha,p}_{n,1} + \frac{8G^{(0,3)}_{\alpha,p,\gamma_n} + 3G^{(0,4)}_{\alpha,p,\gamma_n}}{24n^2},$$

$$\tilde{G}^{\alpha,p}_{n,3} = \tilde{G}^{\alpha,p}_{n,2} - \frac{12G^{(0,4)}_{\alpha,p,\gamma_n} + 8G^{(0,5)}_{\alpha,p,\gamma_n} + G^{(0,6)}_{\alpha,p,\gamma_n}}{48n^3}.$$

4 Proof of Proposition 3.1

Fix $\ell \in \mathbb{N}$ and $\alpha \in (0,1) \cup (1,2)$. The strategy of the proof is the same as the proof of the Proposition of Csörgő (8). It is based on Esseen's classical result (see Petrov (19, Theorem 5.2)).

Lemma 4.1 (Esseen)**.** *Let* F *be a distribution function and* G *be a differentiable function of bounded variation on* $\mathbb{R}$ *with Fourier–Stieltjes transforms* $\mathbf{f}(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\, \mathrm{d}F(x)$ *and* $\mathbf{g}(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\, \mathrm{d}G(x)$, $t \in \mathbb{R}$, *such that* $G(-\infty) := \lim_{x\to-\infty} G(x) = 0$. *Then*

$$\sup_{x\in\mathbb{R}} |F(x) - G(x)| \leq \frac{b}{2\pi} \int_{-T}^{T} \left| \frac{\mathbf{f}(t) - \mathbf{g}(t)}{t} \right| \mathrm{d}t + c_b \frac{\sup_{x\in\mathbb{R}} |G'(x)|}{T}$$

for every choice of $T > 0$ *and* $b > 1$, *where* $c_b > 0$ *is a constant depending only on* b, *which can be given as* $c_b = 4bd_b^2/\pi$, *where* $d_b > 0$ *is the unique root* d *of the equation* $\frac{4}{\pi}\int_0^d \frac{\sin^2 u}{u^2}\, \mathrm{d}u = 1 + \frac{1}{b}$.

We apply this lemma for $F(x) = \mathsf{P}\left\{\frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \le x\right\}$ and $G(x) = G_{n,\ell-1}^{\alpha,p}(x)$, $x \in \mathbb{R}$. By Lemma 6 of Csörgő (8), $G_{n,\ell-1}^{\alpha,p}$ is a differentiable function of bounded variation on $\mathbb{R}$ with $G_{n,\ell-1}^{\alpha,p}(-\infty) = 0$. The Fourier–Stieltjes transform of $G_{n,\ell-1}^{\alpha,p}$ is

$$\mathbf{g}_{n,\ell-1}^{\alpha,p}(t) = \mathrm{e}^{y_{\gamma_n}^{\alpha,p}(t)} \left\{ 1 + \sum_{k=0}^{2\ell-2} \sum_{j=-\lfloor k/2 \rfloor}^{\ell-k-1} \sum_{m=\max\{1,-j\}}^{k+j} \frac{w_{m,k,j}^{\alpha,p} \, (-\mathrm{i}t)^k \, [y_{\gamma_n}^{\alpha,p}(t)]^{j+m}}{m! \, n^{\frac{k}{\alpha}+j}} \right\},$$

for $t \in \mathbb{R}$. The aim of the following discussion is to find an appropriate estimate for $|\mathbf{f}_n^{\alpha,p}(t) - \mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|$ if $|t| \le T_{n,\ell-1}^{\alpha,p}$ with appropriate $T_{n,\ell-1}^{\alpha,p}$. Recalling formula (2.1) for $\mathbf{f}_n^{\alpha,p}$, we have

$$\mathbf{f}_n^{\alpha,p}(t) = \exp\left\{ -\mathrm{i}t\mu_1^{\alpha,p} n^{(\alpha-1)/\alpha} + n \log\left(1 + \frac{x_n^{\alpha,p}(t)}{n}\right) \right\}, \tag{4.1}$$

provided $|x_n^{\alpha,p}(t)| < n$. In order to estimate $|x_n^{\alpha,p}(t)|$, we will use formula (2.2) for $x_n^{\alpha,p}(t)$ containing $y_{\gamma_n}^{\alpha,p}(t)$ and $R_{n,1,2}^{\alpha,p}(t)$. Csörgő (4, Lemma 3), (8, Lemma 3) proved the following crucial estimates.

Lemma 4.2 (Csörgő)**.** *For arbitrary* $\alpha \in (0,1) \cup (1,2)$ *and* $p \in (0,1)$, *there exists* $C_1^{\alpha,p} > 0$ *such that, uniformly in* $\gamma \in (q,1]$,

$$\mathfrak{Re}(y_\gamma^{\alpha,p}(t)) \le -C_1^{\alpha,p}|t|^\alpha, \qquad |y_\gamma^{\alpha,p}(t)| \le C_1^{\alpha,p}|t|^\alpha, \qquad t \in \mathbb{R}.$$

Applying this lemma, we can derive the following estimates.

Lemma 4.3. *For arbitrary* $k \ge 2$, $\alpha \in (0,1) \cup (1,2)$ *and* $p \in (0,1)$, *there exists* $C_{1,k}^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$,

$$|R_{n,1,k}^{\alpha,p}(t)| \le C_{1,k}^{\alpha,p} \frac{|t|^k}{n^{(k-\alpha)/\alpha}}, \qquad t \in \mathbb{R}.$$

Further, for arbitrary $\alpha \in (0,1) \cup (1,2)$ *and* $p \in (0,1)$, *there exist* $C_2^{\alpha,p} > 0$ *and* $C_3^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \le C_2^{\alpha,p} n^{1/\alpha}$,

$$\frac{|x_n^{\alpha,p}(t)|}{n} \le \frac{1}{2}, \qquad |x_n^{\alpha,p}(t)| \le \begin{cases} C_3^{\alpha,p}\,|t|^\alpha, & \text{if } 0 < \alpha < 1, \\ C_3^{\alpha,p}\,|t|\,n^{(\alpha-1)/\alpha}, & \text{if } 1 < \alpha < 2. \end{cases}$$

Proof. By the elementary inequality

$$\left| \mathrm{e}^{\mathrm{i}u} - \sum_{j=0}^{k-1} \frac{(\mathrm{i}u)^j}{j!} \right| \le \frac{|u|^k}{k!}, \qquad u \in \mathbb{R}, \tag{4.2}$$

we obtain

$$\begin{aligned} |R_{n,1,k}^{\alpha,p}(t)| &\le \frac{p\gamma_n}{q} \sum_{m=\lceil \log_r n \rceil}^{\infty} \frac{|t|^k}{k! \, r^{km/\alpha} \, \gamma_n^{k/\alpha}} r^m = \frac{p\,|t|^k}{k!\, q\, \gamma_n^{(k-\alpha)/\alpha}} \frac{q^{\lceil \log_r n \rceil (k-\alpha)/\alpha}}{1 - q^{(k-\alpha)/\alpha}} \\ &= \frac{p}{k!\,(q - q^{k/\alpha})} \frac{|t|^k}{\left(\gamma_n r^{\lceil \log_r n \rceil}\right)^{(k-\alpha)/\alpha}} = -\frac{\mu_k^{\alpha,p}}{k!} \frac{|t|^k}{n^{(k-\alpha)/\alpha}}, \end{aligned}$$

for all $t \in \mathbb{R}$, hence the statement is satisfied with $C_{1,k}^{\alpha,p} = -\mu_k^{\alpha,p}/k! > 0$.

By (2.2), Lemma 4.2 and the estimate for $R_{n,1,2}^{\alpha,p}$,

$$|x_n^{\alpha,p}(t)| \le C_1^{\alpha,p}|t|^\alpha + |\mu_1^{\alpha,p}||t|n^{(\alpha-1)/\alpha} + C_{1,2}^{\alpha,p}\frac{|t|^2}{n^{(2-\alpha)/\alpha}}, \qquad t \in \mathbb{R}, \tag{4.3}$$

hence for $|t| \le C_2^{\alpha,p} n^{1/\alpha}$,

$$\frac{|x_n^{\alpha,p}(t)|}{n} \le C_1^{\alpha,p}(C_2^{\alpha,p})^\alpha + |\mu_1^{\alpha,p}|C_2^{\alpha,p} + C_{1,2}^{\alpha,p}(C_2^{\alpha,p})^2 \le \frac{1}{2}$$

for all sufficiently small $C_2^{\alpha,p} > 0$. If $0 < \alpha < 1$ then, by (4.3), for $|t| \le C_2^{\alpha,p} n^{1/\alpha}$,

$$\begin{aligned} |x_n^{\alpha,p}(t)| &\le |t|^\alpha \left\{C_1^{\alpha,p} + |\mu_1^{\alpha,p}||t|^{1-\alpha}n^{(\alpha-1)/\alpha} + C_{1,2}^{\alpha,p}|t|^{2-\alpha}n^{(\alpha-2)/\alpha}\right\} \\ &\le |t|^\alpha \left\{C_1^{\alpha,p} + |\mu_1^{\alpha,p}|(C_2^{\alpha,p})^{1-\alpha} + C_{1,2}^{\alpha,p}(C_2^{\alpha,p})^{2-\alpha}\right\}. \end{aligned}$$

If $1 < \alpha < 2$ then again by (4.3), for $|t| \le C_2^{\alpha,p} n^{1/\alpha}$,

$$\begin{aligned} |x_n^{\alpha,p}(t)| &\le |t|n^{(\alpha-1)/\alpha} \left\{\frac{C_1^{\alpha,p}|t|^{\alpha-1}}{n^{(\alpha-1)/\alpha}} + |\mu_1^{\alpha,p}| + \frac{C_{1,2}^{\alpha,p}|t|}{n^{1/\alpha}}\right\} \\ &\le |t|n^{(\alpha-1)/\alpha} \left\{C_1^{\alpha,p}(C_2^{\alpha,p})^{\alpha-1} + |\mu_1^{\alpha,p}| + C_{1,2}^{\alpha,p}C_2^{\alpha,p}\right\}, \end{aligned}$$

establishing the case $1 < \alpha < 2$. ☐

Remark that Csörgő (4) also derived these estimates for $|x_n^{\alpha,p}(t)|$.

For $k \in \{0, 1, 2, \dots\}$ and $|t| \le C_2^{\alpha,p} n^{1/\alpha}$, introduce

$$R_{n,2,k}^{\alpha,p}(t) := \sum_{j=k}^{\infty} \frac{1}{j!}\left[R_{n,1,2}^{\alpha,p}(t) + R_{n,3,2}^{\alpha,p}(t)\right]^j, \tag{4.4}$$

where

$$R_{n,3,k}^{\alpha,p}(t) := \sum_{j=k}^{\infty} \frac{(-1)^{j+1}}{j}\frac{[x_n^{\alpha,p}(t)]^j}{n^{j-1}}. \tag{4.5}$$

Then, by Lemma 4.3, for $|t| \le C_2^{\alpha,p} n^{1/\alpha}$, we may write

$$n\log\left(1 + \frac{x_n^{\alpha,p}(t)}{n}\right) = n\sum_{j=1}^{\infty}\frac{(-1)^{j+1}}{j}\left(\frac{x_n^{\alpha,p}(t)}{n}\right)^j = x_n^{\alpha,p}(t) + R_{n,3,2}^{\alpha,p}(t).$$

Next we separate all the terms of order $n^{-\frac{k}{\alpha}-j}$ with $k + j \le \ell - 1$ of the formal expansion of $\mathbf{f}_n^{\alpha,p}$. By (4.1), (2.2) and (2.4), for $|t| \leq C_2^{\alpha,p} n^{1/\alpha}$,

$$\begin{aligned}
\mathbf{f}_n^{\alpha,p}(t) &= \exp\left\{-\mathrm{i}t\mu_1^{\alpha,p} n^{(\alpha-1)/\alpha} + x_n^{\alpha,p}(t) + R_{n,3,2}^{\alpha,p}(t)\right\} \\
&= \exp\left\{y_{\gamma_n}^{\alpha,p}(t) + R_{n,1,2}^{\alpha,p}(t) + R_{n,3,2}^{\alpha,p}(t)\right\} \\
&= \mathrm{e}^{y_{\gamma_n}^{\alpha,p}(t)} \sum_{j=0}^{\infty} \frac{1}{j!} \left[R_{n,1,2}^{\alpha,p}(t) + R_{n,3,2}^{\alpha,p}(t)\right]^j \\
&= \mathrm{e}^{y_{\gamma_n}^{\alpha,p}(t)} \left\{1 + \sum_{j=1}^{\ell-1} \frac{1}{j!} \left[R_{n,1,2}^{\alpha,p}(t) + R_{n,3,2}^{\alpha,p}(t)\right]^j + R_{n,2,\ell}^{\alpha,p}(t)\right\} \\
&= \mathrm{e}^{y_{\gamma_n}^{\alpha,p}(t)} \Bigg\{1 + \sum_{j=1}^{\ell-1} \frac{1}{j!} \Bigg[\sum_{k=2}^{\ell-j+1} \frac{\mu_k^{\alpha,p}\,(\mathrm{i}t)^k}{k!\, n^{\frac{k}{\alpha}-1}} + \sum_{k=2}^{\ell-j+1} \frac{(-1)^{k+1}}{k} \frac{[x_n^{\alpha,p}(t)]^k}{n^{k-1}} + \\
&\qquad\qquad + R_{n,1,\ell-j+2}^{\alpha,p}(t) + R_{n,3,\ell-j+2}^{\alpha,p}(t)\Bigg]^j + R_{n,2,\ell}^{\alpha,p}(t)\Bigg\}.
\end{aligned}$$

Using notation

$$\widetilde{x}_{n,m}^{\alpha,p}(t) := y_{\gamma_n}^{\alpha,p}(t) + \sum_{j=1}^{m} \frac{\mu_j^{\alpha,p}\,(\mathrm{i}t)^j}{j!\, n^{\frac{j}{\alpha}-1}}, \qquad t \in \mathbb{R}, \qquad m \in \mathbb{N}, \tag{4.6}$$

by (2.2), we obtain

$$x_n^{\alpha,p}(t) = \widetilde{x}_{n,m}^{\alpha,p}(t) + R_{n,1,m+1}^{\alpha,p}(t), \qquad t \in \mathbb{R}, \qquad m \in \mathbb{N},$$

and then

$$\mathbf{f}_n^{\alpha,p}(t) = \mathbf{g}_{\alpha,p,\gamma_n}(t) \left\{1 + \sum_{j=1}^{\ell-1} \frac{1}{j!} \left[\widetilde{R}_{n,4,\ell-j+1}^{\alpha,p}(t)\right]^j + R_{n,5,\ell}^{\alpha,p}(t) + R_{n,2,\ell}^{\alpha,p}(t)\right\},$$

where, for $m \in \mathbb{N}$ with $m \geq 2$,

$$\widetilde{R}_{n,4,m}^{\alpha,p}(t) := \sum_{k=2}^{m} \frac{\mu_k^{\alpha,p}\,(\mathrm{i}t)^k}{k!\, n^{\frac{k}{\alpha}-1}} + \sum_{k=2}^{m} \frac{(-1)^{k+1}}{k} \frac{[\widetilde{x}_{n,m-k+1}^{\alpha,p}(t)]^k}{n^{k-1}}, \tag{4.7}$$

$$R_{n,5,\ell}^{\alpha,p}(t) := \sum_{j=1}^{\ell-1} \frac{1}{j!} \sum_{s=1}^{j} \binom{j}{s} \left[R_{n,6,\ell-j+1}^{\alpha,p}(t)\right]^s \left[\widetilde{R}_{n,4,\ell-j+1}^{\alpha,p}(t)\right]^{j-s}, \tag{4.8}$$

with

$$R_{n,6,m}^{\alpha,p}(t) := R_{n,1,m+1}^{\alpha,p}(t) + \widetilde{R}_{n,3,m}^{\alpha,p}(t) + R_{n,3,m+1}^{\alpha,p}(t), \tag{4.9}$$

$$\widetilde{R}_{n,3,m}^{\alpha,p}(t) := \sum_{k=2}^{m} \frac{(-1)^{k+1}}{k\, n^{k-1}} \sum_{u=1}^{k} \binom{k}{u} \left[R_{n,1,m-k+2}^{\alpha,p}(t)\right]^u \left[\widetilde{x}_{n,m-k+1}^{\alpha,p}(t)\right]^{k-u}. \tag{4.10}$$

Clearly $\sum_{j=1}^{\ell-1} \big[\widetilde{R}^{\alpha,p}_{n,4,\ell-j+1}(t)\big]^j/j!$ consists of certain terms of the formal infinite expansion (2.5) of $\mathbf{f}^{\alpha,p}_n(t)$, and all the terms of order $n^{-\frac{k}{\alpha}-j}$ with $k+j \leq \ell-1$ of (2.5) are contained in $\sum_{j=1}^{\ell-1} \big[\widetilde{R}^{\alpha,p}_{n,4,\ell-j+1}(t)\big]^j/j!$. Consequently, for all $|t| \leq C^{\alpha,p}_2 n^{1/\alpha}$,

$$\sum_{j=1}^{\ell-1} \frac{1}{j!}\big[\widetilde{R}^{\alpha,p}_{n,4,\ell-j+1}(t)\big]^j = \sum_{k=0}^{2\ell-2} \sum_{j=-\lfloor k/2 \rfloor}^{\ell-k-1} \sum_{m=\max\{1,-j\}}^{k+j} \frac{w^{\alpha,p}_{m,k,j}\,(-\mathrm{i}t)^k\,[y^{\alpha,p}_{\gamma_n}(t)]^{j+m}}{m!\,n^{\frac{k}{\alpha}+j}} + R^{\alpha,p}_{n,7,\ell}(t), \tag{4.11}$$

where $R^{\alpha,p}_{n,7,\ell}(t)$ contains only terms of order $n^{-\frac{k}{\alpha}-j}$ with $k+j \geq \ell$. Finally, we recognize that

$$\mathbf{f}^{\alpha,p}_n(t) = \mathbf{g}^{\alpha,p}_{n,\ell-1}(t) + \mathrm{e}^{y^{\alpha,p}_{\gamma_n}(t)} R^{\alpha,p}_{n,\ell}(t) \qquad \text{for all } |t| \leq C^{\alpha,p}_2 n^{1/\alpha}, \tag{4.12}$$

where

$$R^{\alpha,p}_{n,\ell}(t) := R^{\alpha,p}_{n,7,\ell}(t) + R^{\alpha,p}_{n,5,\ell}(t) + R^{\alpha,p}_{n,2,\ell}(t).$$

In order to estimate the remainder term $R^{\alpha,p}_{n,\ell}$, we need the following lemmas. Recall the definition (4.5) of $R^{\alpha,p}_{n,3,k}$.

Lemma 4.4. *For all* $k \in \mathbb{N}$, $\alpha \in (0,1) \cup (1,2)$, $p \in (0,1)$ *and* $n \in \mathbb{N}$,

$$|R^{\alpha,p}_{n,3,k}(t)| \leq \frac{2|x^{\alpha,p}_n(t)|^k}{k\,n^{k-1}}, \qquad \textit{for all } |t| \leq C^{\alpha,p}_2 n^{1/\alpha}.$$

Proof. By Lemma 4.3,

$$|R^{\alpha,p}_{n,3,k}(t)| \leq \frac{|x^{\alpha,p}_n(t)|^k}{k\,n^{k-1}} \sum_{j=k}^{\infty} \left|\frac{x^{\alpha,p}_n(t)}{n}\right|^{j-k} \leq \frac{|x^{\alpha,p}_n(t)|^k}{k\,n^{k-1}} \sum_{m=0}^{\infty} \frac{1}{2^m},$$

for all $|t| \leq C^{\alpha,p}_2 n^{1/\alpha}$. □

Lemma 4.5. *For arbitrary* $\ell \in \mathbb{N}$, $\alpha \in (0,1) \cup (1,2)$ *and* $p \in (0,1)$, *there exists* $C^{\alpha,p}_{2,\ell} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \leq C^{\alpha,p}_2 n^{1/\alpha}$,

$$|R^{\alpha,p}_{n,5,\ell}(t)| + |R^{\alpha,p}_{n,7,\ell}(t)| \leq \begin{cases} C^{\alpha,p}_{2,\ell} \dfrac{|t|^{(\ell+1)\alpha} + |t|^{2\ell\alpha}}{n^\ell}, & \textit{if } 0 < \alpha < 1, \\[2ex] C^{\alpha,p}_{2,\ell} \dfrac{|t|^{2+(\ell-1)(2-\alpha)} + |t|^{2\ell}}{n^{\ell(2-\alpha)/\alpha}}, & \textit{if } 1 < \alpha < 2. \end{cases}$$

Proof. A term of order $n^{-\frac{k}{\alpha}-j}$ of $R^{\alpha,p}_{n,7,\ell}(t)$ is contained in the formal infinite expansion (2.5) of $\mathbf{f}^{\alpha,p}_n(t)$, hence it has the form

$$\frac{w^{\alpha,p}_{m,k,j}\,(-\mathrm{i}t)^k\,[y^{\alpha,p}_{\gamma_n}(t)]^{j+m}}{m!\,n^{\frac{k}{\alpha}+j}},$$

where $k \geq 0$, $j \geq -\lfloor k/2 \rfloor$ and $1 \leq m \leq k+j$. By Lemma 4.2, we have

$$\left|\frac{(-\mathrm{i}t)^k\,[y^{\alpha,p}_{\gamma_n}(t)]^{j+m}}{n^{\frac{k}{\alpha}+j}}\right| \leq (C^{\alpha,p}_1)^{j+m} \frac{|t|^{k+(j+m)\alpha}}{n^{\frac{k}{\alpha}+j}}, \qquad n \in \mathbb{N}, \quad t \in \mathbb{R}.$$

For all $n \in \mathbb{N}$ and $|t| \le C_2^{\alpha,p} n^{1/\alpha}$,

$$\frac{|t|^{k+(j+m)\alpha}}{n^{\frac{k}{\alpha}+j}} \le \begin{cases} (C_2^{\alpha,p})^{k(1-\alpha)} \dfrac{|t|^{(k+j+m)\alpha}}{n^{k+j}}, & \text{if } 0<\alpha<1, \\[2ex] (C_2^{\alpha,p})^{(k+2j)(\alpha-1)} \dfrac{|t|^{(k+j)(2-\alpha)+m\alpha}}{n^{(k+j)(2-\alpha)/\alpha}}, & \text{if } 1<\alpha<2. \end{cases}$$

Since $R_{n,7,\ell}^{\alpha,p}(t)$ contains only terms of order $n^{-\frac{k}{\alpha}-j}$ with $k+j \ge \ell$ of the formal infinite expansion (2.5) of $\mathbf{f}_n^{\alpha,p}(t)$, and $1 \le m \le k+j$, we obtain the estimate for $|R_{n,7,\ell}^{\alpha,p}(t)|$. For the estimate of $|R_{n,5,\ell}^{\alpha,p}(t)|$ we derive from Lemmas 4.2, 4.3 and 4.4 the estimates

$$|\widetilde{x}_{n,m}^{\alpha,p}(t)| \le \begin{cases} C_{3,m}|t|^{\alpha}, & \text{if } 0<\alpha<1, \\ C_{3,m}|t| n^{(\alpha-1)/\alpha}, & \text{if } 1<\alpha<2, \end{cases}$$

$$|\widetilde{R}_{n,3,m}^{\alpha,p}(t)| \le \begin{cases} C_{3,m} \dfrac{|t|^{2+(m-1)\alpha}}{n^{\frac{2}{\alpha}+m-2}}, & \text{if } 0<\alpha<1, \\[2ex] C_{3,m} \dfrac{|t|^{m+1}}{n^{\frac{m+1}{\alpha}-1}}, & \text{if } 1<\alpha<2, \end{cases}$$

$$|\widetilde{R}_{n,4,m}^{\alpha,p}(t)| \le \begin{cases} C_{3,m} \dfrac{|t|^{2\alpha}}{n}, & \text{if } 0<\alpha<1, \\[2ex] C_{3,m} \dfrac{|t|^{2}}{n^{(2-\alpha)/\alpha}}, & \text{if } 1<\alpha<2, \end{cases}$$

$$|R_{n,6,m}^{\alpha,p}(t)| \le \begin{cases} C_{3,m} \dfrac{|t|^{(m+1)\alpha}}{n^{m}}, & \text{if } 0<\alpha<1, \\[2ex] C_{3,m} \dfrac{|t|^{m+1}}{n^{\frac{m+1}{\alpha}-1}}, & \text{if } 1<\alpha<2, \end{cases}$$

for all $n \in \mathbb{N}$ and $|t| \le C_2^{\alpha,p} n^{1/\alpha}$ and with sufficiently large constants $C_{3,m} > 0$. The last two inequlaities imply the estimate for $|R_{n,5,\ell}^{\alpha,p}(t)|$. $\square$

Lemma 4.6. *For arbitrary* $\ell \in \mathbb{N}$, $\alpha \in (0,1) \cup (1,2)$ *and* $p \in (0,1)$, *there exist* $C_{4,\ell}^{\alpha,p} > 0$ *and* $\varepsilon_{\alpha,p} \in \big(0, C_2^{\alpha,p}\big]$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \le \varepsilon_{\alpha,p} n^{1/\alpha}$,

$$|R_{n,2,\ell}^{\alpha,p}(t)| \le \begin{cases} C_{4,\ell}^{\alpha,p} \dfrac{|t|^{2\ell\alpha} \, \mathrm{e}^{C_1^{\alpha,p}|t|^{\alpha}/2}}{n^{\ell}}, & \textit{if } 0<\alpha<1, \\[2ex] C_{4,\ell}^{\alpha,p} \dfrac{|t|^{2\ell} \, \mathrm{e}^{C_1^{\alpha,p}|t|^{\alpha}/2}}{n^{\ell(2-\alpha)/\alpha}}, & \textit{if } 1<\alpha<2. \end{cases}$$

Proof. Recalling the definition (4.4) of $R^{\alpha,p}_{n,2,\ell}$, for all $n \in \mathbb{N}$ and $|t| \leq C^{\alpha,p}_2 n^{1/\alpha}$,

$$\begin{aligned} |R^{\alpha,p}_{n,2,\ell}(t)| &\leq \sum_{j=\ell}^{\infty} \frac{1}{j!} \big(|R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)|\big)^j \\ &\leq \frac{\big(|R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)|\big)^\ell}{\ell!} \sum_{j=\ell}^{\infty} \frac{\big(|R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)|\big)^{j-\ell}}{(j-\ell)!} \qquad (4.13) \\ &= \frac{\big(|R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)|\big)^\ell}{\ell!} \mathrm{e}^{|R^{\alpha,p}_{n,1,2}(t)|+|R^{\alpha,p}_{n,3,2}(t)|}. \end{aligned}$$

By Lemmas 4.3 and 4.4, for all $n \in \mathbb{N}$ and $|t| \leq C^{\alpha,p}_2 n^{1/\alpha}$,

$$|R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)| \leq C^{\alpha,p}_{1,2} \frac{t^2}{n^{(2-\alpha)/\alpha}} + \frac{|x^{\alpha,p}_n(t)|^2}{n}.$$

If $\alpha \in (0,1)$ then by Lemma 4.3, for all $n \in \mathbb{N}$ and $|t| \leq \varepsilon_{\alpha,p} n^{1/\alpha}$ with $\varepsilon_{\alpha,p} \in \big(0, C^{\alpha,p}_2\big]$,

$$\begin{aligned} |R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)| &\leq C^{\alpha,p}_{1,2} \frac{t^2}{n^{(2-\alpha)/\alpha}} + \frac{(C^{\alpha,p}_3)^2\,|t|^{2\alpha}}{n} \\ &\leq \big[C^{\alpha,p}_{1,2}(\varepsilon_{\alpha,p})^{2-2\alpha} + (C^{\alpha,p}_3)^2\big] \frac{|t|^{2\alpha}}{n} \qquad (4.14) \\ &\leq \big[C^{\alpha,p}_{1,2}(\varepsilon_{\alpha,p})^{2-2\alpha} + (C^{\alpha,p}_3)^2\big] (\varepsilon_{\alpha,p})^\alpha\,|t|^\alpha \leq \frac{1}{2} C^{\alpha,p}_1\,|t|^\alpha \qquad (4.15) \end{aligned}$$

for sufficiently small $\varepsilon_{\alpha,p} > 0$. Applying in the inequality (4.13) the estimates (4.14) and (4.15) for $\big(|R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)|\big)^\ell$ and $\mathrm{e}^{|R^{\alpha,p}_{n,1,2}(t)|+|R^{\alpha,p}_{n,3,2}(t)|}$, respectively, we obtain the statement for $\alpha \in (0,1)$.

If $\alpha \in (1,2)$ then for all $n \in \mathbb{N}$ and $|t| \leq \varepsilon_{\alpha,p} n^{1/\alpha}$ with $\varepsilon_{\alpha,p} \in \big(0, C^{\alpha,p}_2\big]$,

$$\begin{aligned} |R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)| &\leq \big[C^{\alpha,p}_{1,2} + (C^{\alpha,p}_3)^2\big] \frac{t^2}{n^{(2-\alpha)/\alpha}} \qquad (4.16) \\ &\leq \big[C^{\alpha,p}_{1,2} + (C^{\alpha,p}_3)^2\big] (\varepsilon_{\alpha,p})^{2-\alpha}\,|t|^\alpha \leq \frac{1}{2} C^{\alpha,p}_1\,|t|^\alpha \qquad (4.17) \end{aligned}$$

for sufficiently small $\varepsilon_{\alpha,p} > 0$. Applying in the inequality (4.13) the estimates (4.16) and (4.17) for $\big(|R^{\alpha,p}_{n,1,2}(t)| + |R^{\alpha,p}_{n,3,2}(t)|\big)^\ell$ and $\mathrm{e}^{|R^{\alpha,p}_{n,1,2}(t)|+|R^{\alpha,p}_{n,3,2}(t)|}$, respectively, we obtain the statement for $\alpha \in (1,2)$. □

By (4.12), the first inequality of Lemma 4.2, and Lemmas 4.5 and 4.6, we obtain

Lemma 4.7. *For arbitrary* $\ell \in \mathbb{N}$, $\alpha \in (0,1) \cup (1,2)$ *and* $p \in (0,1)$, *there exists* $C^{\alpha,p}_{5,\ell} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \leq \varepsilon_{\alpha,p} n^{1/\alpha}$,

$$|\mathbf{f}^{\alpha,p}_n(t) - \mathbf{g}^{\alpha,p}_{n,\ell-1}(t)| \leq \begin{cases} C^{\alpha,p}_{5,\ell} \dfrac{|t|^{(\ell+1)\alpha} + |t|^{2\ell\alpha}}{n^\ell} \mathrm{e}^{-C^{\alpha,p}_1 |t|^\alpha/2}, & \textit{if } \alpha \in (0,1), \\[2ex] C^{\alpha,p}_{5,\ell} \dfrac{|t|^{2+(\ell-1)(2-\alpha)} + |t|^{2\ell}}{n^{\ell(2-\alpha)/\alpha}} \mathrm{e}^{-C^{\alpha,p}_1 |t|^\alpha/2}, & \textit{if } \alpha \in (1,2). \end{cases}$$

Now by Lemma 4.1,

$$\Delta_{n,\ell}^{\alpha,p} := \sup_{x\in\mathbb{R}} \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \leq x \right\} - G_{n,\ell-1}^{\alpha,p}(x) \right| \leq \frac{b}{\pi} \Delta_{n,\ell,1}^{\alpha,p} + c_b \Delta_{n,\ell,2}^{\alpha,p}$$

for every $b > 1$, where

$$\Delta_{n,\ell,1}^{\alpha,p} := \int_0^{\varepsilon_{\alpha,p}\, n^{1/\alpha}} \frac{|\mathbf{f}_n^{\alpha,p}(t) - \mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|}{|t|} \,\mathrm{d}t,$$

$$\Delta_{n,\ell,2}^{\alpha,p} := \frac{M_{n,\ell}^{\alpha,p}}{\varepsilon_{\alpha,p}\, n^{1/\alpha}} \qquad \text{with} \quad M_{n,\ell}^{\alpha,p} := \sup_{x\in\mathbb{R}} \left| \frac{\mathrm{d}G_{n,\ell-1}^{\alpha,p}(x)}{\mathrm{d}x} \right|.$$

By Lemmas 4 and 6 of Csörgő (8), $\sup_{n\in\mathbb{N}} M_{n,\ell}^{\alpha,p} < \infty$, and hence $\Delta_{n,\ell,2}^{\alpha,p} = O\left(\frac{1}{n^{1/\alpha}}\right)$ for all fixed $\ell \in \mathbb{N}$. Using the simple fact that

$$\int_0^\infty t^\beta \mathrm{e}^{-Ct^\alpha} \,\mathrm{d}t < \infty, \qquad \beta > -1, \qquad C > 0, \tag{4.18}$$

by Lemma 4.7 we obtain

$$\Delta_{n,\ell,1}^{\alpha,p} = O\left(\frac{1}{n^\ell} + \frac{1}{n^{\ell(2-\alpha)/\alpha}} \right). \tag{4.19}$$

Consequently, we conclude $\Delta_{n,\ell}^{\alpha,p} = O\left(\frac{1}{n^\ell} + \frac{1}{n^{\ell(2-\alpha)/\alpha}} + \frac{1}{n^{1/\alpha}}\right)$, which implies the first statement.

The reduction of the order $O\left(\frac{1}{n^{1/\alpha}}\right)$ to $o\left(\frac{1}{n^{1/\alpha}}\right)$ when $\ell \in \{2, 3, \dots\}$, $\alpha \in \left(\frac{1}{\ell}, 1\right) \cup \left(1, 2 - \frac{1}{\ell}\right)$ and $r^{1/\alpha} \notin \mathbb{N}$ is based on the following classical result due to Esseen (10).

Lemma 4.8 (Esseen). *If* $\mathbf{f}$ *is the characteristic function of a non-lattice distribution, then for every fixed* $\varepsilon > 0$, *there exists a sequence* $\lambda_n \to \infty$ *such that*

$$\int_\varepsilon^{\lambda_n} \frac{|\mathbf{f}(t)|^n}{t} \,\mathrm{d}t = o\left(\mathrm{e}^{-\sqrt{n}/2}\right) \qquad \textit{as} \ \ n \to \infty.$$

It is easy to check that the distribution of X is non-lattice if and only if $r^{1/\alpha} \notin \mathbb{N}$. Thus there exists a sequence $\lambda_n^{\alpha,p} \to \infty$ such that

$$\begin{aligned} I_n^{\alpha,p} &:= \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\lambda_n^{\alpha,p}\, n^{1/\alpha}} \frac{|\mathbf{f}_n^{\alpha,p}(t)|}{t} \,\mathrm{d}t = \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\lambda_n^{\alpha,p}\, n^{1/\alpha}} \frac{|\mathbf{f}_{\alpha,p}(t/n^{1/\alpha})|^n}{t} \,\mathrm{d}t \\ &= \int_{\varepsilon_{\alpha,p}}^{\lambda_n^{\alpha,p}} \frac{|\mathbf{f}_{\alpha,p}(s)|^n}{s} \,\mathrm{d}s = o\left(\mathrm{e}^{-\sqrt{n}/2}\right) = o\left(\frac{1}{n^{1/\alpha}}\right) \qquad \text{as} \ \ n \to \infty. \end{aligned}$$

Clearly (4.19) implies $\Delta_{n,\ell,1}^{\alpha,p} = o\left(\frac{1}{n^{1/\alpha}}\right)$ as $n \to \infty$ for $\alpha \in \left(\frac{1}{\ell}, 1\right) \cup \left(1, 2 - \frac{1}{\ell}\right)$. Hence by Lemma 4.1 now with $T = \lambda_n^{\alpha,p}\, n^{1/\alpha}$ we have

$$\Delta_{n,\ell}^{\alpha,p} \leq \frac{b}{\pi} \left(\Delta_{n,\ell,1}^{\alpha,p} + I_n^{\alpha,p} + I_{n,\ell}^{\alpha,p} \right) + c_b \frac{M_{n,\ell}^{\alpha,p}}{\lambda_n^{\alpha,p} n^{1/\alpha}} = \frac{b}{\pi} I_{n,\ell}^{\alpha,p} + o\left(\frac{1}{n^{1/\alpha}}\right)$$

as $n \to \infty$, where

$$I^{\alpha,p}_{n,\ell} := \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\infty} \frac{|\mathbf{g}^{\alpha,p}_{n,\ell-1}(t)|}{t}\, \mathrm{d}t = O\left(n^{\ell-2} \mathrm{e}^{-C_1^{\alpha,p}(\varepsilon_{\alpha,p})^{\alpha}\, n}\right) = o\left(\frac{1}{n^{1/\alpha}}\right)$$

as $n \to \infty$, using Lemma 4.2 and the simple fact that

$$\int_n^{\infty} u^{\beta} \mathrm{e}^{-u}\, \mathrm{d}u = O(n^{\beta}\mathrm{e}^{-n}), \qquad \beta \in \mathbb{R}. \tag{4.20}$$

Thus $\Delta^{\alpha,p}_{n,\ell} = o\left(\frac{1}{n^{1/\alpha}}\right)$ as $n \to \infty$ for $\alpha \in \left(\frac{1}{\ell}, 1\right) \cup \left(1, 2 - \frac{1}{\ell}\right)$.

5 Nonuniform bounds in asymptotic expansions

The main results are contained in the following

Proposition 5.1. *For* $\ell \in \{2, 3, \dots\}$ *and* $\alpha \in \left(1, 2 - \frac{1}{\ell}\right]$,

$$\sup_{x \in \mathbb{R}} (1 + |x|) \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \le x \right\} - G^{\alpha,p}_{n,\ell-1}(x) \right| = O\left(\frac{1}{n^{1/\alpha}}\right).$$

For $\ell \in \mathbb{N}$ *and* $\alpha \in \left(2 - \frac{1}{\ell},\, 2\right)$,

$$\sup_{x \in \mathbb{R}} (1 + |x|) \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \le x \right\} - G^{\alpha,p}_{n,\ell-1}(x) \right| = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right).$$

For $\ell \in \{2, 3, \dots\}$, $\alpha \in \left(1,\, 2 - \frac{1}{\ell}\right)$ *and* $r^{1/\alpha} \notin \mathbb{N}$,

$$\sup_{x \in \mathbb{R}} (1 + |x|) \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \le x \right\} - G^{\alpha,p}_{n,\ell-1}(x) \right| = o\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right).$$

For $\ell = 1$ and $\alpha \in (1, 2)$, we have a nonuniform rate of merge

$$\sup_{x \in \mathbb{R}} (1 + |x|) \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \le x \right\} - G_{\alpha,p,\gamma_n}(x) \right| = O\left(\frac{1}{n^{(2-\alpha)/\alpha}}\right).$$

Certain terms of $G^{\alpha,p}_{n,\ell-1}$ are of the same or of a smaller order than the remainder terms $O\left(\frac{1}{n^{1/\alpha}}\right)$ or $o\left(\frac{1}{n^{1/\alpha}}\right)$. Using the boundedness of the functions $G^{(k,j)}_{\alpha,p,\gamma_n}$, the expansions may be simplified as follows. Recall the definition of the approximation functions $\tilde{G}^{\alpha,p}_{n,\ell}$ and $\tilde{\tilde{G}}^{\alpha,p}_{n,\ell}$ from Theorem 3.1.

Theorem 5.1. *For* $\ell \in \{2, 3, \dots\}$ *and* $\alpha \in \left(2 - \frac{1}{\ell-1},\, 2 - \frac{1}{\ell}\right]$,

$$\sup_{x \in \mathbb{R}} (1 + |x|) \left| \mathsf{P}\left\{ \frac{S_n - c_n^{\alpha,p}}{n^{1/\alpha}} \le x \right\} - \tilde{G}^{\alpha,p}_{n,\ell-1}(x) \right| = O\left(\frac{1}{n^{1/\alpha}}\right).$$

For $\ell = 2$, $\alpha \in \left(1, \frac{3}{2}\right)$, *or for* $\ell \in \{3, 4, \dots\}$, $\alpha \in \left[2 - \frac{1}{\ell-1},\, 2 - \frac{1}{\ell}\right)$, *and for* $r^{1/\alpha} \notin \mathbb{N}$,

$$\sup_{x \in \mathbb{R}} (1 + |x|) \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \le x \right\} - \tilde{\tilde{G}}^{\alpha,p}_{n,\ell-1}(x) \right| = o\left(\frac{1}{n^{1/\alpha}}\right),$$

6 Proof of Proposition 5.1

Fix $\ell \in \mathbb{N}$ and $\alpha \in (1,2)$. The proof is based on a result due to Osipov (17) (see Petrov (19, Lemmas 6.7, 6.8) and Hall (11, Lemma 1.1); note that the result has been stated only for $k \geq 2$, but the proof works for $k = 1$ as well).

Lemma 6.1 (Osipov)**.** *Let F be a distribution function and G be a differentiable function of bounded variation on $\mathbb{R}$ with Fourier–Stieltjes transforms $\mathbf{f}(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\,\mathrm{d}F(x)$ and $\mathbf{g}(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\,\mathrm{d}G(x)$, $t \in \mathbb{R}$, such that $G(-\infty) = \lim_{x\to-\infty} G(x) = 0$, $G(+\infty) = \lim_{x\to+\infty} G(x) = 1$, and*

$$\int_{-\infty}^{\infty} |x|^k \,|\mathrm{d}(F(x) - G(x))| < \infty$$

with some $k \in \mathbb{N}$. Then the function $H_k(x) := x^k(F(x) - G(x))$, $x \in \mathbb{R}$, is of bounded variation on $\mathbb{R}$ with Fourier–Stieltjes transform

$$\mathbf{h}_k(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\,\mathrm{d}H_k(x) = \frac{k!}{(-\mathrm{i}t)^k} \sum_{j=0}^{k} \frac{(-t)^j}{j!} \frac{\mathrm{d}^j(\mathbf{f}-\mathbf{g})(t)}{\mathrm{d}t^j},$$

for $t \in \mathbb{R}$ with $t \neq 0$, and there exists $c_k > 0$ such that

$$\begin{aligned} &\sup_{x\in\mathbb{R}} (1 + |x|^k)|F(x) - G(x)| \\ &\leq c_k \int_{-T}^{T} \left| \frac{\mathbf{f}(t) - \mathbf{g}(t)}{t} \right| \mathrm{d}t + c_k \int_{-T}^{T} \left| \frac{\mathbf{h}_k(t)}{t} \right| \mathrm{d}t + c_k \frac{\sup_{x\in\mathbb{R}}(1+|x|^k)|G'(x)|}{T} \end{aligned}$$

for every choice of $T > 0$.

We apply this lemma for $F(x) = F_n^{\alpha,p}(x) = \mathsf{P}\left\{\frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \leq x\right\}$ and $G(x) = G_{n,\ell-1}^{\alpha,p}(x)$, $x \in \mathbb{R}$, introduced in Section 3. By Lemma 6 of Csörgő (8), $G_{n,\ell-1}^{\alpha,p}$ is a differentiable function of bounded variation on $\mathbb{R}$ with $G_{n,\ell-1}^{\alpha,p}(-\infty) = 0$ and $G_{n,\ell-1}^{\alpha,p}(+\infty) = 1$. Since $\alpha \in (1,2)$, the expectation $\mathsf{E}(X)$ of the gain in one game is finite, which implies $\int_{-\infty}^{\infty} |x|\,\mathrm{d}F_n^{\alpha,p}(x) = \mathsf{E}\left|\frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}}\right| < \infty$. By Lemma 6 of Csörgő (8), $\alpha \in (1,2)$ also implies $\int_{-\infty}^{\infty} |x|\,|\mathrm{d}G_{n,\ell-1}^{\alpha,p}(x)| < \infty$, since $G_{n,\ell-1}^{\alpha,p}$ is a linear combination of the continuously differentiable functions $G_{\alpha,p,\gamma}^{(k,j)}$, $k, j \in \{0,1,2,\ldots\}$, $\gamma \in (q,1]$, and $\int_{-\infty}^{\infty} |x|\,|\mathrm{d}G_{\alpha,p,\gamma}^{(k,j)}(x)| = \int_{-\infty}^{\infty} |x||G_{\alpha,p,\gamma}^{(k+1,j)}(x)|\,\mathrm{d}x < \infty$. Consequently,

$$\int_{-\infty}^{\infty} |x|\,|\mathrm{d}(F_n^{\alpha,p}(x) - G_{n,\ell-1}^{\alpha,p}(x))| \leq \int_{-\infty}^{\infty} |x|\,\mathrm{d}F_n^{\alpha,p}(x) + \int_{-\infty}^{\infty} |x|\,|G_{n,\ell-1}^{\alpha,p}(x))| < \infty.$$

By Lemma 6.1, the function $H_{n,\ell-1}^{\alpha,p}(x) := x(F_n^{\alpha,p}(x) - G_{n,\ell-1}^{\alpha,p}(x))$, $x \in \mathbb{R}$, is of bounded variation on $\mathbb{R}$ with Fourier–Stieltjes transform

$$\mathbf{h}_{n,\ell-1}^{\alpha,p}(t) = \int_{-\infty}^{\infty} \mathrm{e}^{\mathrm{i}tx}\,\mathrm{d}H_{n,\ell-1}^{\alpha,p}(x) = \frac{\mathrm{i}}{t}(\mathbf{f}_n^{\alpha,p}(t) - \mathbf{g}_{n,\ell-1}^{\alpha,p}(t)) - \mathrm{i}\frac{\mathrm{d}(\mathbf{f}_n^{\alpha,p} - \mathbf{g}_{n,\ell-1}^{\alpha,p})(t)}{\mathrm{d}t}$$

for $t \in \mathbb{R}$ with $t \neq 0$. Now we apply Lemma 6.1 for $T = \varepsilon_{\alpha,p} n^{1/\alpha}$ with $\varepsilon_{\alpha,p} > 0$ from Lemmas 4.6 and 4.7, and we obtain

$$\begin{aligned}
\widetilde{\Delta}_{n,\ell}^{\alpha,p} &:= \sup_{x\in\mathbb{R}} (1+|x|) \left| \mathsf{P}\left\{ \frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \leq x \right\} - G_{n,\ell-1}^{\alpha,p}(x) \right| \\
&\leq 2c_1 \Big(\widetilde{\Delta}_{n,\ell,1}^{\alpha,p} + \widetilde{\Delta}_{n,\ell,2}^{\alpha,p} + \widetilde{\Delta}_{n,\ell,3}^{\alpha,p} \Big) + c_1 \widetilde{\Delta}_{n,\ell,4}^{\alpha,p},
\end{aligned}$$

where

$$\begin{aligned}
\widetilde{\Delta}_{n,\ell,1}^{\alpha,p} &:= \int_0^{\varepsilon_{\alpha,p} n^{1/\alpha}} \frac{|\mathbf{f}_n^{\alpha,p}(t) - \mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|}{|t|} \, \mathrm{d}t, \\
\widetilde{\Delta}_{n,\ell,2}^{\alpha,p} &:= \int_0^{\varepsilon_{\alpha,p} n^{1/\alpha}} \frac{|\mathbf{f}_n^{\alpha,p}(t) - \mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|}{|t|^2} \, \mathrm{d}t, \\
\widetilde{\Delta}_{n,\ell,3}^{\alpha,p} &:= \int_0^{\varepsilon_{\alpha,p} n^{1/\alpha}} \frac{1}{|t|} \left| \frac{\mathrm{d}(\mathbf{f}_n^{\alpha,p} - \mathbf{g}_{n,\ell-1}^{\alpha,p})(t)}{\mathrm{d}t} \right| \mathrm{d}t, \\
\widetilde{\Delta}_{n,\ell,4}^{\alpha,p} &:= \frac{\widetilde{M}_{n,\ell}^{\alpha,p}}{\varepsilon_{\alpha,p} n^{1/\alpha}} \qquad \text{with} \quad \widetilde{M}_{n,\ell}^{\alpha,p} := \sup_{x\in\mathbb{R}} (1+|x|) \left| \frac{\mathrm{d}G_{n,\ell-1}^{\alpha,p}(x)}{\mathrm{d}x} \right|.
\end{aligned}$$

By Lemmas 4 and 6 of Csörgő (8), $\sup_{n\in\mathbb{N}} \widetilde{M}_{n,\ell}^{\alpha,p} < \infty$, and hence $\widetilde{\Delta}_{n,\ell,4}^{\alpha,p} = O\left(\frac{1}{n^{1/\alpha}}\right)$ for all fixed $\ell \in \mathbb{N}$. By (4.19), we have $\widetilde{\Delta}_{n,\ell,1}^{\alpha,p} = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right)$. By (4.12) and Lemmas 4.2, 4.5 and 4.6 we obtain

$$\widetilde{\Delta}_{n,\ell,2}^{\alpha,p} \leq \int_0^{\varepsilon_{\alpha,p} n^{1/\alpha}} \frac{|R_{n,\ell}^{\alpha,p}(t)| \, \mathrm{e}^{-C_1^{\alpha,p}|t|^\alpha}}{|t|^2} \, \mathrm{d}t = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}} \right).$$

The aim of the following discussion is to find an appropriate estimate for $\left| \frac{\mathrm{d}(\mathbf{f}_n^{\alpha,p} - \mathbf{g}_{n,\ell-1}^{\alpha,p})(t)}{\mathrm{d}t} \right|$ if $|t| \leq \varepsilon_{\alpha,p} n^{1/\alpha}$. Using formula (4.12), first we calculate the derivatives of the ingredients of $\mathbf{f}_n^{\alpha,p} - \mathbf{g}_{n,\ell-1}^{\alpha,p}$. The function $y_\gamma^{\alpha,p}$ is differentiable and

$$\frac{\mathrm{d}y_\gamma^{\alpha,p}(t)}{\mathrm{d}t} = \sum_{k=-\infty}^{\infty} \left(\exp\left\{ \frac{\mathrm{i}tr^{k/\alpha}}{\gamma^{1/\alpha}} \right\} - 1 \right) \frac{\mathrm{i}p\gamma^{(\alpha-1)/\alpha}}{qr^{k(\alpha-1)/\alpha}}, \qquad t \in \mathbb{R}, \tag{6.1}$$

since $\alpha \in (1,2)$ implies absolute convergence of this series. Differentiability of $y_\gamma^{\alpha,p}$ implies that for each $m \in \mathbb{N}$, the function $\widetilde{x}_{n,m}^{\alpha,p}$, introduced in (4.6), hence, for each $m \geq 2$, the function $\widetilde{R}_{n,4,m}^{\alpha,p}$, given in (4.7), and hence for each $k \in \mathbb{N}$, the function $R_{n,7,k}^{\alpha,p}$, defined in (4.11), is differentiable on the whole $\mathbb{R}$. For each $k \geq 2$, the function $R_{n,1,k}^{\alpha,p}$, given in (2.3), is differentiable and

$$\frac{\mathrm{d}R_{n,1,k}^{\alpha,p}(t)}{\mathrm{d}t} = -\sum_{m=\lceil \log_r n \rceil}^{\infty} \left(\exp\left\{ \frac{\mathrm{i}t}{r^{m/\alpha}\gamma_n^{1/\alpha}} \right\} - \sum_{j=0}^{k-2} \frac{(\mathrm{i}t)^j}{j!\, r^{jm/\alpha}\gamma_n^{j/\alpha}} \right) \frac{\mathrm{i}p\gamma_n^{(\alpha-1)/\alpha}}{q} r^{m(\alpha-1)/\alpha}, \tag{6.2}$$

for $t \in \mathbb{R}$, since $\alpha \in (1,2)$ implies absolute convergence of this series. Differentiability of $\widetilde{x}_{n,m}^{\alpha,p}$, $m \in \mathbb{N}$, and $R_{n,1,k}^{\alpha,p}$, $k \geq 2$, imply that for each $m \geq 2$, the function $\widetilde{R}_{n,3,m}^{\alpha,p}$, given in

(4.10), is differentiable on the whole $\mathbb{R}$. Recalling formula (2.2), we obtain differentiability of the function $x_n^{\alpha,p}$, and

$$\frac{\mathrm{d}x_n^{\alpha,p}(t)}{\mathrm{d}t} = \frac{\mathrm{d}y_{\gamma_n}^{\alpha,p}(t)}{\mathrm{d}t} + \mathrm{i}\mu_1^{\alpha,p} n^{(\alpha-1)/\alpha} + \frac{\mathrm{d}R_{n,1,2}^{\alpha,p}(t)}{\mathrm{d}t}, \qquad t \in \mathbb{R}. \tag{6.3}$$

For each $k \in \mathbb{N}$, the function $R_{n,3,k}^{\alpha,p}$, defined in (4.5), is differentiable for $|t| < C_2^{\alpha,p} n^{1/\alpha}$ and

$$\frac{\mathrm{d}R_{n,3,k}^{\alpha,p}(t)}{\mathrm{d}t} = \frac{\mathrm{d}x_n^{\alpha,p}(t)}{\mathrm{d}t} \sum_{j=k-1}^{\infty} \frac{(-1)^j [x_n^{\alpha,p}(t)]^j}{n^j}, \qquad |t| < C_2^{\alpha,p} n^{1/\alpha}, \tag{6.4}$$

since by Lemma 4.3, $\alpha \in (1,2)$ implies absolute convergence of this series for $|t| < C_2^{\alpha,p} n^{1/\alpha}$. For each $k \in \mathbb{N}$, the function $R_{n,2,k}^{\alpha,p}$, given in (4.4), is differentiable for $|t| < C_2^{\alpha,p} n^{1/\alpha}$ and

$$\frac{\mathrm{d}R_{n,2,k}^{\alpha,p}(t)}{\mathrm{d}t} = \left(\frac{\mathrm{d}R_{n,1,2}^{\alpha,p}(t)}{\mathrm{d}t} + \frac{\mathrm{d}R_{n,3,2}^{\alpha,p}(t)}{\mathrm{d}t} \right) R_{n,2,k-1}^{\alpha,p}(t), \tag{6.5}$$

for $|t| < C_2^{\alpha,p} n^{1/\alpha}$. Differentiability of $R_{n,1,k}^{\alpha,p}$, $k \geq 2$, $\widetilde{R}_{n,3,m}^{\alpha,p}$, $m \in \mathbb{N}$, and $R_{n,3,k}^{\alpha,p}$, $k \in \mathbb{N}$, imply that for each $m \geq 2$, the function $R_{n,6,m}^{\alpha,p}$, given in (4.9), and hence for each $k \in \mathbb{N}$, the function $R_{n,5,k}^{\alpha,p}$, introduced in (4.8), is differentiable for $|t| < C_2^{\alpha,p} n^{1/\alpha}$. Consequently by (4.12), we conclude

$$\begin{aligned} \frac{\mathrm{d}(\mathbf{f}_n^{\alpha,p} - \mathbf{g}_{n,\ell-1}^{\alpha,p})(t)}{\mathrm{d}t} &= \mathrm{e}^{y_{\gamma_n}^{\alpha,p}(t)} \left(\frac{\mathrm{d}R_{n,2,\ell}^{\alpha,p}(t)}{\mathrm{d}t} + \frac{\mathrm{d}R_{n,5,\ell}^{\alpha,p}(t)}{\mathrm{d}t} + \frac{\mathrm{d}R_{n,7,\ell}^{\alpha,p}(t)}{\mathrm{d}t} \right) \\ &\quad + \frac{\mathrm{d}y_{\gamma_n}^{\alpha,p}(t)}{\mathrm{d}t} \big(\mathbf{f}_n^{\alpha,p}(t) - \mathbf{g}_{n,\ell-1}^{\alpha,p}(t) \big), \qquad |t| < C_2^{\alpha,p} n^{1/\alpha}. \end{aligned}$$

In order to estimate $\left| \frac{\mathrm{d}(\mathbf{f}_n^{\alpha,p} - \mathbf{g}_{n,\ell-1}^{\alpha,p})(t)}{\mathrm{d}t} \right|$, we need the following lemmas.

Lemma 6.2. *For arbitrary* $\alpha \in (1,2)$ *and* $p \in (0,1)$, *there exists* $C_4^{\alpha,p} > 0$ *such that, uniformly in* $\gamma \in (q,1]$,

$$\mathfrak{Im} \left(\frac{\mathrm{d}y_\gamma^{\alpha,p}(t)}{\mathrm{d}t} \right) \leq -C_4^{\alpha,p} |t|^{\alpha-1}, \qquad \left| \frac{\mathrm{d}y_\gamma^{\alpha,p}(t)}{\mathrm{d}t} \right| \leq C_4^{\alpha,p} |t|^{\alpha-1}, \qquad t \in \mathbb{R}.$$

Proof. These estimates can be derived in the same way as the inequality (4) in the proof of Lemma 3 of Csörgő (4) and the second statement of Lemma 3 in Csörgő (8), using (6.1). □

The following lemmas can be proved as Lemmas 4.3, 4.4, 4.5, 4.6 and 4.7, respectively.

Lemma 6.3. *For arbitrary* $k \geq 2$, $\alpha \in (1,2)$ *and* $p \in (0,1)$, *there exists* $C_{5,k}^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$,

$$\left| \frac{\mathrm{d}R_{n,1,k}^{\alpha,p}(t)}{\mathrm{d}t} \right| \leq C_{5,k}^{\alpha,p} \frac{|t|^{k-1}}{n^{(k-\alpha)/\alpha}}, \qquad t \in \mathbb{R}.$$

Further, for arbitrary $\alpha \in (1,2)$ *and* $p \in (0,1)$, *there exists* $C_5^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \leq C_2^{\alpha,p} n^{1/\alpha}$,

$$\left| \frac{\mathrm{d}x_n^{\alpha,p}(t)}{\mathrm{d}t} \right| \leq C_5^{\alpha,p} n^{(\alpha-1)/\alpha}.$$

Lemma 6.4. *For arbitrary* $k \geq 2$, $\alpha \in (1,2)$ *and* $p \in (0,1)$, *there exists* $C_{6,k}^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \leq C_2^{\alpha,p}\, n^{1/\alpha}$,

$$\left|\frac{\mathrm{d}R_{n,3,k}^{\alpha,p}(t)}{\mathrm{d}t}\right| \leq C_{6,k}^{\alpha,p} \frac{|t|^{k-1}}{n^{(k-\alpha)/\alpha}}.$$

Lemma 6.5. *For arbitrary* $\ell \in \mathbb{N}$, $\alpha \in (1,2)$ *and* $p \in (0,1)$, *there exists* $C_{7,\ell}^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \leq C_2^{\alpha,p}\, n^{1/\alpha}$,

$$\left|\frac{\mathrm{d}R_{n,5,\ell}^{\alpha,p}(t)}{\mathrm{d}t}\right| + \left|\frac{\mathrm{d}R_{n,7,\ell}^{\alpha,p}(t)}{\mathrm{d}t}\right| \leq C_{7,\ell}^{\alpha,p} \frac{|t|^{1+(\ell-1)(2-\alpha)} + |t|^{2\ell-1}}{n^{\ell(2-\alpha)/\alpha}}.$$

Lemma 6.6. *For arbitrary* $\ell \in \mathbb{N}$, $\alpha \in (1,2)$ *and* $p \in (0,1)$, *there exists* $C_{8,\ell}^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \leq \varepsilon_{\alpha,p}\, n^{1/\alpha}$,

$$\left|\frac{\mathrm{d}R_{n,2,\ell}^{\alpha,p}(t)}{\mathrm{d}t}\right| \leq C_{8,\ell}^{\alpha,p} \frac{|t|^{2\ell-1}\, \mathrm{e}^{C_1^{\alpha,p}\,|t|^\alpha/2}}{n^{\ell(2-\alpha)/\alpha}}.$$

Lemma 6.7. *For arbitrary* $\ell \in \mathbb{N}$, $\alpha \in (0,1) \cup (1,2)$ *and* $p \in (0,1)$, *there exist* $C_{9,\ell}^{\alpha,p} > 0$ *such that for all* $n \in \mathbb{N}$ *and* $|t| \leq \varepsilon_{\alpha,p}\, n^{1/\alpha}$,

$$\left|\frac{\mathrm{d}(\mathbf{f}_n^{\alpha,p} - \mathbf{g}_{n,\ell-1}^{\alpha,p})(t)}{\mathrm{d}t}\right| \leq C_{9,\ell}^{\alpha,p} \frac{|t|^{1+(\ell-1)(2-\alpha)} + |t|^{2\ell-1+\alpha}}{n^{\ell(2-\alpha)/\alpha}} \mathrm{e}^{-C_1^{\alpha,p}\,|t|^\alpha/2}.$$

By Lemma 6.7 and the inequality (4.18), we obtain $\widetilde{\Delta}_{n,\ell,3}^{\alpha,p} = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right)$. Consequently, we conclude $\widetilde{\Delta}_{n,\ell}^{\alpha,p} = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}} + \frac{1}{n^{1/\alpha}}\right)$, which implies the first statement.

The reduction of the order $O\left(\frac{1}{n^{1/\alpha}}\right)$ to $o\left(\frac{1}{n^{1/\alpha}}\right)$ when $\ell \in \{2,3,\dots\}$, $\alpha \in \left(1, 2-\frac{1}{\ell}\right)$ and $r^{1/\alpha} \notin \mathbb{N}$ is based again on Lemma 4.8. By this lemma, there exists a sequence $\lambda_n^{\alpha,p} \to \infty$ such that

$$\int_{\varepsilon_{\alpha,p}}^{\lambda_n^{\alpha,p}} \frac{|\mathbf{f}_{\alpha,p}(s)|^n}{s}\,\mathrm{d}s = o\left(\mathrm{e}^{-\sqrt{n}/2}\right) \qquad \text{as } n \to \infty.$$

Then

$$\begin{aligned} \tilde{I}_{n,1}^{\alpha,p} &:= \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\lambda_{n-1}^{\alpha,p}\, n^{1/\alpha}} \frac{|\mathbf{f}_n^{\alpha,p}(t)|}{t}\,\mathrm{d}t = \int_{\varepsilon_{\alpha,p}}^{\lambda_{n-1}^{\alpha,p}} \frac{|\mathbf{f}_{\alpha,p}(s)|^n}{s}\,\mathrm{d}s \\ &\leq \int_{\varepsilon_{\alpha,p}}^{\lambda_{n-1}^{\alpha,p}} \frac{|\mathbf{f}_{\alpha,p}(s)|^{n-1}}{s}\,\mathrm{d}s = o\left(\mathrm{e}^{-\sqrt{n-1}/2}\right) = o\left(\frac{1}{n^{1/\alpha}}\right) \qquad \text{as } n \to \infty, \end{aligned}$$

and similarly,

$$\tilde{I}_{n,2}^{\alpha,p} := \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\lambda_{n-1}^{\alpha,p}\, n^{1/\alpha}} \frac{|\mathbf{f}_n^{\alpha,p}(t)|}{t^2}\,\mathrm{d}t = o\left(n^{-1/\alpha}\,\mathrm{e}^{-\sqrt{n-1}/2}\right) = o\left(\frac{1}{n^{1/\alpha}}\right),$$

as $n \to \infty$. By (2.1),

$$\frac{\mathrm{d}\mathbf{f}_n^{\alpha,p}(t)}{\mathrm{d}t} = n^{(\alpha-1)/\alpha}\big(\mathbf{f}_{\alpha,p}(t/n^{1/\alpha})\big)^{n-1}\,\mathbf{f}'_{\alpha,p}(t/n^{1/\alpha})\,\mathrm{e}^{-\mathrm{i}t\mu_1^{\alpha,p}n^{(\alpha-1)/\alpha}} - \mathrm{i}\,\mu_1^{\alpha,p}\,n^{(\alpha-1)/\alpha}\,\mathbf{f}_n^{\alpha,p}(t). \tag{6.6}$$

It is easy to check that $\mathbf{f}'_{\alpha,p}(t) = \mathrm{i}\,\mathsf{E}(X\mathrm{e}^{\mathrm{i}tX})$ for all $t \in \mathbb{R}$, thus $|\mathbf{f}'_{\alpha,p}(t)| \leq \mu_1^{\alpha,p}$ for all $t \in \mathbb{R}$, hence

$$\tilde{I}_{n,3}^{\alpha,p} := \int_{\varepsilon_{\alpha,p} n^{1/\alpha}}^{\lambda_{n-1} n^{1/\alpha}} \frac{1}{t} \left| \frac{\mathrm{d}\mathbf{f}_n^{\alpha,p}(t)}{\mathrm{d}t} \right| \mathrm{d}t = o\left(n^{(\alpha-1)/\alpha}\, \mathrm{e}^{-\sqrt{n-1}/2} \right) = o\left(\frac{1}{n^{1/\alpha}} \right),$$

as $n \to \infty$. By Lemma 6.1 now with $T = \lambda_{n-1}^{\alpha,p}\, n^{1/\alpha}$,

$$\widetilde{\Delta}_{n,\ell}^{\alpha,p} \leq 2c_1 \sum_{j=1}^{3} \left(\widetilde{\Delta}_{n,\ell,j}^{\alpha,p} + \tilde{I}_{n,j}^{\alpha,p} \right) + 2c_1 \tilde{\tilde{I}}_{n,\ell}^{\alpha,p} + c_1 \frac{\widetilde{M}_{n,\ell}^{\alpha,p}}{\lambda_{n-1}^{\alpha,p}\, n^{1/\alpha}} \qquad \text{as } n \to \infty,$$

where, by the first inequality of Lemma 4.2, formula (6.1), and the statement (4.20),

$$\begin{aligned} \tilde{\tilde{I}}_{n,\ell}^{\alpha,p} &:= \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\infty} \left(\frac{|\mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|}{t} + \frac{|\mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|}{t^2} + \frac{1}{t} \left| \frac{\mathrm{d}\mathbf{g}_{n,\ell-1}^{\alpha,p}(t)}{\mathrm{d}t} \right| \right) \mathrm{d}t \\ &= O\left(n^{\ell-2-\frac{1}{\alpha}}\, \mathrm{e}^{-C_1^{\alpha,p}(\varepsilon_{\alpha,p})^{\alpha} n} \right) = o\left(\frac{1}{n^{1/\alpha}} \right), \qquad \text{as } n \to \infty. \end{aligned}$$

Thus $\widetilde{\Delta}_{n,\ell}^{\alpha,p} = o\left(\frac{1}{n^{1/\alpha}}\right)$ as $n \to \infty$ for $\alpha \in \left(1, 2 - \frac{1}{\ell}\right)$.

7 Uniform and nonuniform bounds in asymptotic expansions in local merging theorems in the lattice case

Theorem 7.1. *For* $\ell \in \mathbb{N}$ *and* $r^{1/\alpha} \in \mathbb{N}$,

$$\sup_{s \in r^{1/\alpha}\mathbb{N}} \left| \frac{n^{1/\alpha}}{r^{1/\alpha}}\, \mathsf{P}\{S_n = s\} - \left(G_{n,\ell-1}^{\alpha,p}\right)' \left(\frac{s - c_n^{\alpha,p}}{n^{1/\alpha}} \right) \right| = \begin{cases} O\left(\frac{1}{n^{\ell}}\right), & \text{if } 0 < \alpha < 1, \\ O\left(\frac{[\log_r n]^{2\ell}}{n^{\ell}}\right), & \text{if } \alpha = 1, \\ O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right), & \text{if } 1 < \alpha < 2. \end{cases}$$

In case $\ell = 1$ Theorem 7.1 implies for all $0 < \alpha < 2$ the local merging theorem

$$\lim_{n \to \infty} \left[\frac{n^{1/\alpha}}{r^{1/\alpha}}\, \mathsf{P}\{S_n = s\} - G'_{\alpha,p,\gamma_n} \left(\frac{s - c_n^{\alpha,p}}{n^{1/\alpha}} \right) \right] = 0$$

for all $s \in r^{1/\alpha}\mathbb{N}$.

Theorem 7.2. *For* $\ell \in \mathbb{N}$, $\alpha \in (1,2)$ *and* $r^{1/\alpha} \in \mathbb{N}$,

$$\sup_{s \in r^{1/\alpha}\mathbb{N}} \left(1 + \frac{|s - \mu_1^{\alpha,p} n|}{n^{1/\alpha}} \right) \left| \frac{n^{1/\alpha}}{r^{1/\alpha}}\, \mathsf{P}\{S_n = s\} - \left(G_{n,\ell-1}^{\alpha,p}\right)' \left(\frac{s - \mu_1^{\alpha,p} n}{n^{1/\alpha}} \right) \right| = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}} \right).$$

8 Proof Theorems 7.1 and 7.2

First we prove Theorem 7.1. Fix $\ell \in \mathbb{N}$ and $\alpha \in (0,2)$ such that $r^{1/\alpha} \in \mathbb{N}$. We have

$$\mathbf{f}_n^{\alpha,p}(t) = \mathsf{E}\left(\mathrm{e}^{\mathrm{i}t(S_n - c_n^{\alpha,p})/n^{1/\alpha}}\right) = \sum_{u \in r^{1/\alpha}\mathbb{N}} \mathrm{e}^{\mathrm{i}t(u - c_n^{\alpha,p})/n^{1/\alpha}}\, \mathsf{P}\{S_n = u\},$$

for all $t \in \mathbb{R}$, hence for every $s \in r^{1/\alpha}\mathbb{N}$,

$$\int_{-\pi n^{1/\alpha}/r^{1/\alpha}}^{\pi n^{1/\alpha}/r^{1/\alpha}} \mathrm{e}^{-\mathrm{i}t(s - c_n^{\alpha,p})/n^{1/\alpha}} \mathbf{f}_n^{\alpha,p}(t)\,\mathrm{d}t$$

$$= \sum_{u \in r^{1/\alpha}\mathbb{N}} \mathsf{P}\{S_n = u\} \int_{-\pi n^{1/\alpha}/r^{1/\alpha}}^{\pi n^{1/\alpha}/r^{1/\alpha}} \mathrm{e}^{\mathrm{i}t(u-s)/n^{1/\alpha}}\,\mathrm{d}t = \frac{2\pi n^{1/\alpha}}{r^{1/\alpha}}\,\mathsf{P}\{S_n = s\}.$$

(In fact, this is the inversion formula for probabilities.) By Lemma 4 in Csörgő (8),

$$\left(G_{n,\ell-1}^{\alpha,p}\right)'(x) = \frac{1}{2\pi}\int_{-\infty}^{\infty} \mathrm{e}^{-\mathrm{i}tx} \mathbf{g}_{n,\ell-1}^{\alpha,p}(t)\,\mathrm{d}t, \qquad x \in \mathbb{R}.$$

Thus we have

$$D_{n,\ell}^{\alpha,p} := \sup_{s \in r^{1/\alpha}\mathbb{N}} \left| \frac{n^{1/\alpha}}{r^{1/\alpha}}\,\mathsf{P}\{S_n = s\} - \left(G_{n,\ell-1}^{\alpha,p}\right)'\left(\frac{s - c_n^{\alpha,p}}{n^{1/\alpha}}\right)\right| \leq \frac{1}{\pi}\left(D_{n,\ell,1}^{\alpha,p} + J_n^{\alpha,p} + J_{n,\ell}^{\alpha,p}\right),$$

where

$$D_{n,\ell,1}^{\alpha,p} := \int_0^{\varepsilon_{\alpha,p}\, n^{1/\alpha}} |\mathbf{f}_n^{\alpha,p}(t) - \mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|\,\mathrm{d}t,$$

$$J_n^{\alpha,p} := \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\pi n^{1/\alpha}/r^{1/\alpha}} |\mathbf{f}_n^{\alpha,p}(t)|\,\mathrm{d}t,$$

$$J_{n,\ell}^{\alpha,p} := \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\infty} |\mathbf{g}_{n,\ell-1}^{\alpha,p}(t)|\,\mathrm{d}t,$$

with $\varepsilon_{\alpha,p} > 0$ from Lemma 4.6.

First consider the case $\alpha \in (0,1) \cup (1,2)$. Then by Lemma 4.7,

$$D_{n,\ell,1}^{\alpha,p} = O\left(\frac{1}{n^\ell} + \frac{1}{n^{\ell(2-\alpha)/\alpha}}\right).$$

Moreover,

$$J_n^{\alpha,p} = \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\pi n^{1/\alpha}/r^{1/\alpha}} \left|\mathbf{f}_{\alpha,p}\left(\frac{t}{n^{1/\alpha}}\right)\right|^n \mathrm{d}t = n^{1/\alpha}\int_{\varepsilon_{\alpha,p}}^{\pi/r^{1/\alpha}} |\mathbf{f}_{\alpha,p}(t)|^n\,\mathrm{d}t$$

$$\leq \frac{\pi n^{1/\alpha}}{r^{1/\alpha}}\left(\sup_{t \in [\varepsilon_{\alpha,p},\, \pi/r^{1/\alpha}]} |\mathbf{f}_{\alpha,p}(t)|\right)^n = O\left(\frac{1}{n^\ell} + \frac{1}{n^{\ell(2-\alpha)/\alpha}}\right),$$

since

$$\sup_{t\in[\varepsilon_{\alpha,p},\,\pi/r^{1/\alpha}]} |\mathbf{f}_{\alpha,p}(t)| < 1. \tag{8.1}$$

(This follows from the fact that the gain X in one game takes values also in the lattice $r^{1/\alpha}\mathbb{Z}$, the maximal span of a lattice L with $\mathsf{P}\{X \in L\} = 1$ is $r^{1/\alpha}$, hence $\sup_{t\in H} |\,\mathsf{E}(\mathrm{e}^{\mathrm{i}tX})| < 1$ for all compact set $H \subset \mathbb{R}$ with $H \cap \frac{2\pi}{r^{1/\alpha}}\mathbb{Z} = \emptyset$, see, e.g., Bhattacharya and Ranga Rao (1, §21).)

By Lemma 4.2,

$$J^{\alpha,p}_{n,\ell} = O\left(n^{\ell-2}\,\mathrm{e}^{-C_1^{\alpha,p}(\varepsilon_{\alpha,p})^\alpha\, n}\right) = O\left(\frac{1}{n^\ell} + \frac{1}{n^{\ell(2-\alpha)/\alpha}}\right),$$

and we obtain the statement for $\alpha \in (0,1) \cup (1,2)$.

In case $\alpha = 1$, in order to estimate $D^{1,p}_{n,\ell,1}$, we need an analogue of Lemma 4.7.

Lemma 8.1. *For arbitrary* $\ell \in \mathbb{N}$ *and* $p \in (0,1)$*, there exist* $C^{1,p}_{5,\ell} > 0$ *and* $\varepsilon_{1,p} > 0$*, such that for all* $n \in \mathbb{N}$ *and* $|t| \le \varepsilon_{1,p}\, n$*,*

$$|\mathbf{f}^{1,p}_n(t) - \mathbf{g}^{1,p}_{n,\ell-1}(t)| \le C^{1,p}_{5,\ell}\frac{|t|^{\ell+1} + |t|^{2\ell}}{n^\ell}\left(1 + \log_r \frac{2n}{|t|}\right)^{2\ell} \mathrm{e}^{-C_1^{1,p}|t|/2}.$$

Proof. It can be derived in a similar way as Lemma 4.7. First recall that there exists $C_1^{1,p} > 0$ such that, uniformly in $\gamma \in (q,1]$, we have $|y^{1,p}_\gamma(t)| \le C_1^{1,p}(1 + \log_r|t|)\,|t|$ for all $t \in \mathbb{R}$, see Csörgő (8, Lemma 3). There exists $C_1^{1,p} > 0$ such that, uniformly in $\gamma \in (q,1]$, we have $\mathfrak{Re}(y^{1,p}_\gamma(t)) \le -C_1^{1,p}|t|$ for all $t \in \mathbb{R}$, see inequality (4) in the proof of Lemma 3 of Csörgő (4). Next one can show, as in Lemma 4.3, that there exist $C_2^{1,p} > 0$ and $C_3^{1,p} > 0$ such that for all $n \in \mathbb{N}$ and $|t| \le C_2^{1,p} n$,

$$\frac{|x^{1,p}_n(t)|}{n} \le \frac{1}{2}, \qquad |x^{1,p}_n(t)| \le C_3^{1,p}|t|\left(1 + \log_r \frac{2n}{|t|}\right).$$

The expression for $\mathbf{f}^{1,p}_n(t)$ if $|t| \le C_2^{1,p} n$ has the same form as in case $\alpha \in (0,1) \cup (1,2)$ by replacing α by 1 and $\mu_1^{1,p}$ by $pr\,\log_r n$, hence

$$\widetilde{x}^{1,p}_{n,m}(t) := y^{1,p}_{\gamma_n}(t) + \mathrm{i}tpr\,\log_r n + \sum_{j=2}^{m} \frac{\mu_j^{1,p}\,(\mathrm{i}t)^j}{j!\,n^{j-1}}.$$

Lemma 4.4 is valid for $\alpha = 1$ as well. As in Lemma 4.5, there exists $C^{1,p}_{2,\ell} > 0$ such that for all $n \in \mathbb{N}$ and $|t| \le C_2^{1,p} n$,

$$|R^{1,p}_{n,5,\ell}(t)| + |R^{1,p}_{n,7,\ell}(t)| \le C^{1,p}_{2,\ell}\frac{|t|^{\ell+1} + |t|^{2\ell}}{n^\ell}\left(1 + \log_r \frac{2n}{|t|}\right)^{2\ell}.$$

As in Lemma 4.6, there exist $C_{4,\ell}^{1,p} > 0$ and $\varepsilon_{1,p} \in (0, C_2^{1,p}]$ such that for all $n \in \mathbb{N}$ and $|t| \leq \varepsilon_{1,p}\, n$,

$$|R_{n,2,\ell}^{1,p}(t)| \leq C_{4,\ell}^{1,p} \frac{|t|^{2\ell}}{n^\ell} \left(1 + \log_r \frac{2n}{|t|}\right)^{2\ell} \mathrm{e}^{C_1^{1,p}|t|/2},$$

and we obtain the statement of the lemma. $\square$

Now by Lemma 8.1, we obtain $D_{n,\ell}^{1,p} = O\left(\frac{[\log_r n]^{2\ell}}{n^\ell}\right)$. Clearly, $J_n^{1,p} = O\left(\frac{[\log_r n]^{2\ell}}{n^\ell}\right)$ can be proved as in case $\alpha \in (0,1) \cup (1,2)$. Consequently, $J_{n,\ell}^{1,p} = O\left(\frac{[\log_r n]^{2\ell}}{n^\ell}\right)$, and we conclude the statement of Theorem 7.1 for $\alpha = 1$.

In order to prove Theorem 7.2, first we recall that $\alpha \in (1,2)$ implies $\mathsf{E}(X) < \infty$, and hence $\mathsf{E}\left|\frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}}\right| < \infty$, and the characteristic function $\mathbf{f}_n^{\alpha,p}$ of $\frac{S_n - \mu_1^{\alpha,p} n}{n^{1/\alpha}}$ is differentiable and

$$\frac{\mathrm{d}\mathbf{f}_n^{\alpha,p}(t)}{\mathrm{d}t} = \sum_{u \in r^{1/\alpha}\mathbb{N}} \frac{\mathrm{i}(u - \mu_1^{\alpha,p} n)}{n^{1/\alpha}} \mathrm{e}^{\mathrm{i}t(u - \mu_1^{\alpha,p} n)/n^{1/\alpha}}\, \mathsf{P}\{S_n = u\}, \qquad t \in \mathbb{R}.$$

Consequently, for every $s \in r^{1/\alpha}\mathbb{N}$,

$$\begin{aligned}
&\int_{-\pi n^{1/\alpha}/r^{1/\alpha}}^{\pi n^{1/\alpha}/r^{1/\alpha}} \mathrm{e}^{-\mathrm{i}t(s - \mu_1^{\alpha,p} n)/n^{1/\alpha}} \left(\frac{\mathrm{d}\mathbf{f}_n^{\alpha,p}(t)}{\mathrm{d}t}\right) \mathrm{d}t \\
&\qquad = \sum_{u \in r^{1/\alpha}\mathbb{N}} \frac{\mathrm{i}(u - \mu_1^{\alpha,p} n)}{n^{1/\alpha}}\, \mathsf{P}\{S_n = u\} \int_{-\pi n^{1/\alpha}/r^{1/\alpha}}^{\pi n^{1/\alpha}/r^{1/\alpha}} \mathrm{e}^{\mathrm{i}t(u-s)/n^{1/\alpha}}\, \mathrm{d}t \\
&\qquad = \frac{\mathrm{i}(s - \mu_1^{\alpha,p} n)}{n^{1/\alpha}} \frac{2\pi n^{1/\alpha}}{r^{1/\alpha}}\, \mathsf{P}\{S_n = s\}.
\end{aligned}$$

By Lemmas 4 and 6 in Csörgő (8), $\alpha \in (1,2)$ implies

$$\mathrm{i}x\big(G_{n,\ell-1}^{\alpha,p}\big)'(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \mathrm{e}^{-\mathrm{i}tx} \left(\frac{\mathrm{d}\mathbf{g}_{n,\ell-1}^{\alpha,p}(t)}{\mathrm{d}t}\right) \mathrm{d}t, \qquad x \in \mathbb{R}.$$

Thus we have

$$\begin{aligned}
\widetilde{D}_{n,\ell}^{\alpha,p} &:= \sup_{s \in r^{1/\alpha}\mathbb{N}} \frac{|s - \mu_1^{\alpha,p} n|}{n^{1/\alpha}} \left| \frac{n^{1/\alpha}}{r^{1/\alpha}}\, \mathsf{P}\{S_n = s\} - \big(G_{n,\ell-1}^{\alpha,p}\big)'\Big(\frac{s - \mu_1^{\alpha,p} n}{n^{1/\alpha}}\Big) \right| \\
&\leq \frac{1}{\pi} \left(\widetilde{D}_{n,\ell,1}^{\alpha,p} + \tilde{J}_n^{\alpha,p} + \tilde{J}_{n,\ell}^{\alpha,p}\right),
\end{aligned}$$

where

$$\widetilde{D}_{n,\ell,1}^{\alpha,p} := \int_0^{\varepsilon_{\alpha,p}\, n^{1/\alpha}} \left| \frac{\mathrm{d}\mathbf{f}_n^{\alpha,p}(t)}{\mathrm{d}t} - \frac{\mathrm{d}\mathbf{g}_{n,\ell-1}^{\alpha,p}(t)}{\mathrm{d}t} \right| \mathrm{d}t,$$

$$\tilde{J}_n^{\alpha,p} := \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\pi n^{1/\alpha}/r^{1/\alpha}} \left| \frac{\mathrm{d}\mathbf{f}_n^{\alpha,p}(t)}{\mathrm{d}t} \right| \mathrm{d}t,$$

$$\tilde{J}_{n,\ell}^{\alpha,p} := \int_{\varepsilon_{\alpha,p}\, n^{1/\alpha}}^{\infty} \left| \frac{\mathrm{d}\mathbf{g}_{n,\ell-1}^{\alpha,p}(t)}{\mathrm{d}t} \right| \mathrm{d}t.$$

By Lemma 6.7 and inequality (4.18), $\widetilde{D}_{n,\ell,1}^{\alpha,p} = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right)$. Using (6.6) and the inequalities (8.1) and $|\mathbf{f}'_{\alpha,p}(t)| \leq \mu_1^{\alpha,p}$, $t \in \mathbb{R}$, we obtain $\tilde{J}_n^{\alpha,p} = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right)$. By Lemma 6.2, $\tilde{J}_{n,\ell}^{\alpha,p} = O\left(\frac{1}{n^{\ell(2-\alpha)/\alpha}}\right)$, and by Theorem 7.1, we conclude the statement of Theorem 7.2.

Acknowledgements

I am thanksful to Professor Sándor Csörgő who explained me the phenomena of merging in generalized St. Petersburg games. This research has been supported by the Hungarian Scientific Research Funds under Grant No. T048544 and T079128.

References

[1] Bhattacharya, R. N. and Ranga Rao, R. (1976). *Normal Approximation and Asymptotic Expansions.* Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, New York-London-Sydney. MR0436272

[2] Bikelis, A. (1966). Estimates of the remainder term in the central limit theorem. *Litovsk. Mat. Sb.* **6(3)** 323–346. MR0210173

[3] Christoph, G. and Wolf, W. (1992). *Convergence Theorems with a Stable Limit Law.* Mathematical Research 70, Akademie Verlag, Berlin. MR1202035

[4] Csörgő, S. (2002). Rates of merge in generalized St. Petersburg games. *Acta Sci. Math. (Szeged)* **68** 815–847. MR1954550

[5] Csörgő, S. (2003). Merge rates for sums of large gains in generalized St. Petersburg games. *Acta Sci. Math. (Szeged)* **69** 441–454. MR1992319

[6] Csörgő, S. (2005). A probabilistic proof of Kruglov's theorem on tails of infinitely divisible distributions. *Acta Sci. Math. (Szeged)* **71** 405–415. MR2160375

[7] Csörgő, S. (2007). Fourier analysis of semistable distributions. *Acta Appl. Math.* **96** 159–175. MR2327532

[8] Csörgő, S. (2007). Merging asymptotic expansions in generalized St. Petersburg games. *Acta Sci. Math. (Szeged)* **73** 297–331.

[9] Csörgő, S. and Dodunekova, R. (1991). Limit theorems for the Petersburg game. In: *Sums, Trimmed Sums and Extremes*, Progress in Probability 23, pp. 285–315, Boston, Birkhäuser. MR1117274

[10] Esseen, C.-G. (1945). Fourier analysis of distribution functions. A mathematical study of the Laplace–Gaussian law. *Acta Math.* **77** 1–125. MR0014626

[11] Hall, P. (1982). *Rates of Convergence in the Central Limit Theorem.* Research Notes in Mathematics 62, Pitman, Boston. MR0668197

[12] Hall, P. (1983). Fast rates of convergence in the central limit theorem. *Z. Wahrsch. Verw. Gebiete* **62** 491–507. MR0690574

[13] Katznelson, Y. (1968). *An Introduction to Harmonic Analysis.* John Wiley & Sons, Inc., New York-London-Sydney. MR0248482

[14] Kevei, P. (2009). Merging asymptotic expansions for semistable random variables, *Lithuanian Math. J.* **49(1)** 40–54.

[15] Martin-Löf, A. (1985). A limit theorem which clarifies the 'Petersburg paradox. *J. Appl. Probab.* **22** 634–643. MR0799286

[16] Osipov, L. V. (1967). Asymptotic expansions in the central limit theorem. *Vestnik Leningrad. Univ.* **1967(19)** 45–62. MR0216552

[17] Osipov, L. V. (1972). Asymptotic expansions of the distribution function of a sum of random variables with non uniform estimates for the remainder term. *Vestnik Leningrad. Univ.* **1972(1)** 51–59. MR0300324

[18] Osipov, L. V. and Petrov, V. V. (1967). On the estimation of the remainder term in the central limit theorem. *Teor. Verojatnost. i Primenen.* **12** 322–329. MR0216552

[19] Petrov, V. V. (1975). *Sums of Independent Random Variables.* Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 82. Springer-Verlag, New York-Heidelberg. MR0388499